

- (1) Find a parameterization of the surface formed by the graph of $z = x^2 - y^2$ with (x, y) in the triangle in the xy -plane formed by the x -axis, the y -axis, and the line $y = -x + 1$.
- (2) Is the surface in the previous problem a smooth surface? If no, at what points is it not smooth?
- (3) Find a parameterization of the surface formed by rotating the curve $\begin{pmatrix} \cos t + 5 \\ 2 \sin t \end{pmatrix}$ with $0 \leq t \leq 2\pi$ around the y -axis.
- (4) Consider the surface

$$\mathbf{X}(s, t) = \begin{pmatrix} 2 \sin 3t + t \\ \cos 2s \\ t^2 + s^2 \end{pmatrix}, \quad 0 \leq t \leq \pi/4, \quad 0 \leq s \leq \pi$$

Find the tangent and normal vectors to \mathbf{X} at the point $(\pi/6, \pi/6)$. Is the surface smooth?

- (5) Suppose that $\mathbf{F}: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is a C^1 vector field, and that $\mathbf{X}: D \rightarrow \mathbb{R}^3$ is a smooth, oriented surface. Let $h: E \rightarrow D$ be a smooth, orientation reversing change-of coordinate function. Prove that

$$\iint_{\mathbf{X}} \mathbf{F} \cdot d\mathbf{S} = - \iint_{\mathbf{X} \circ h} \mathbf{F} \cdot d\mathbf{S}.$$

- (6) Suppose that $f: \mathbb{R}^3 \rightarrow \mathbb{R}$ is a C^1 vector field, and that $\mathbf{X}: D \rightarrow \mathbb{R}^3$ is a smooth, oriented surface. Let $h: E \rightarrow D$ be a smooth change-of coordinate function. Prove that


$$\iint_{\mathbf{X}} f \, dS = \iint_{\mathbf{X} \circ h} f \, dS.$$

- (7) Suppose that $\mathbf{X}: D \rightarrow \mathbb{R}^3$ is a smooth, oriented surface with unit normal \mathbf{n} . Suppose that $\mathbf{F}: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is a C^1 vector field. Prove that

$$\iint_{\mathbf{X}} \mathbf{F} \cdot d\mathbf{S} = \iint_{\mathbf{X}} \mathbf{F} \cdot \mathbf{n} \, dS.$$

- (8) Use the previous result to integrate the vector field $\mathbf{F}(x, y, z) = (x, y, z)$ over the unit sphere (with outward normal) in \mathbb{R}^3 .

- (9) Let S be the disc of radius 1 centered at $(1,0,0)$ in \mathbb{R}^3 which is parallel to the yz -plane. Orient S with normal vector pointing in the direction of the positive x -axis. Use the definition of surface integral to calculate the flux of $\mathbf{F}(x,y,z) = (-xy, yz, xz)$ through S .
- (10) Use the same surface S and \mathbf{F} as in the previous problem, but now use Stoke's theorem to calculate the flux of the curl of the previous problem.
- (11) Let S be a surface formed by rotating the image of $\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} t \\ \sin t \end{pmatrix}$, $2\pi \leq t \leq 3\pi$ around the y -axis. Orient S so that at the point $(2\pi + \pi/2, 1, 0)$ there is an upward pointing normal vector. For the following vector fields, find the flux of the vector field through S . (Hint: there are easy ways and there are hard ways...)



(a) $\mathbf{F}(x,y,z) = \begin{pmatrix} x+y \\ -y+z \\ -x+y \end{pmatrix}$

(b) $\mathbf{F}(x,y,z) = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$

(c) $\mathbf{F}(x,y,z) = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$

(d) $\mathbf{F}(x,y,z) = \frac{1}{x^2+z^2} \begin{pmatrix} -z \\ 0 \\ x \end{pmatrix}$

(12) Give a complete, precise statement of Stokes' theorem