

Partial Differential Equations

Some Supplementary notes for MA411

Evan Randles

Spring 2023

Contents

1	Some Calculus of Complex-Valued Functions	1
1.1	Differential Calculus	1
1.2	Integral Calculus	3
2	Sturm-Liouville Theory	4
3	Life on the circle	7
3.1	Integration on \mathbb{T} and some important spaces of functions on \mathbb{T}	8
3.2	Pointwise and Uniform convergence of functions on \mathbb{T}	11
3.3	The Lebesgue norms on \mathbb{T} and the $L^2(\mathbb{T})$ inner product	13
3.4	Fourier coefficients and the uniform theory	15
3.5	Convolutions	21
4	All things Fourier	21
4.1	The L^2 theory	21
4.2	The pointwise and uniform theory theory	28
4.3	The Gibb's phenomenon	34
5	The Fourier Transform	38
A	Sups, Infs and all that	38
B	The Riemann-Darboux integral	42
C	The essence of convergence	51
C.1	Properties of Uniform Convergence	58
C.2	The Weierstrass M -test	60
C.3	Defining Convergence with the Integral: A glimpse at Lebesgue norms	62

These are some supplementary course notes for Partial Differential Equations (Math 411). Please check these frequently as I will often make corrections and changes without explicit warning. Also, if you find or suspect an error typo – no matter how trivial – please email me to let me know!

1 Some Calculus of Complex-Valued Functions

Let $I \subseteq \mathbb{R}$ be an interval, i.e., $I = [a, b]$, (a, b) , $[a, b)$ or $(a, b]$ for extended¹ real numbers $a < b$. Most of the time, $I = \mathbb{R}$, $[0, 1]$, or $[-\pi, \pi]$. In these notes, we shall work with complex-valued functions $f : I \rightarrow \mathbb{C}$ which are functions of the form

$$f(x) = u(x) + iv(x)$$

where u and v are real-valued functions on I . We call u and v the real and imaginary parts of f , respectively, and sometimes write $u = \operatorname{Re}(f)$ and $v = \operatorname{Im}(f)$. We say that a complex-valued function f is real-valued if $v = \operatorname{Im}(f)$ is identically 0 on I . Everything done below generalizes the single-variable calculus you learned long ago (where everything was real-valued) and, in fact, recaptures it exactly when the complex-valued functions have everywhere vanishing imaginary parts.

1.1 Differential Calculus

We shall focus mostly on those complex-valued functions $f : I \rightarrow \mathbb{C}$ which are continuous, i.e., complex-valued functions $f = u + iv$ whose real and imaginary parts u and v are continuous on the interval I . The set of such functions is denoted by $C^0(I)$ and explicitly defined by

$$C^0(I) = \{f : I \rightarrow \mathbb{C} : u = \operatorname{Re}(f) \text{ and } v = \operatorname{Im}(f) \text{ are continuous real-valued functions on } I\}$$

A function $f = u + iv$ is said to be differentiable on I provided that

$$u'(x) = \lim_{h \rightarrow 0} \frac{u(x+h) - u(x)}{h} \quad \text{and} \quad v'(x) = \lim_{h \rightarrow 0} \frac{v(x+h) - v(x)}{h}$$

exist at each² $x \in I$. If we know that f is differentiable on I , we define its derivative f' by

$$\frac{d}{dx} f = f'(x) = u'(x) + iv'(x)$$

for $x \in [0, 1]$. When it happens that f' is itself a continuous function on I , i.e., $f' \in C^0(I)$, we say that f is continuously differentiable³ on I and we denote the set of all such functions by $C^1(I)$, i.e.,

$$C^1(I) = \{f \in C^0(I) : f \text{ is differentiable on } I \text{ and } f' \in C^0(I)\}.$$

Continuing inductively, for each $k \geq 2$, we say that f is k -times differentiable on I provided that the $(k-1)$ th derivative of f is itself differentiable on I and we write

$$\frac{d^k f}{dx^k} = f^{(k)}(x) = \frac{d}{dx} f^{(k-1)}(x) = \frac{d^k}{dx^k} u(x) + \frac{d^k}{dx^k} v(x)$$

for $x \in I$. The set of k -times continuously differentiable functions is defined by

$$C^k(I) = \{f \in C^{k-1}(I) : f^{(k)} \text{ exists and is a member of } C^0(I)\}.$$

¹By extended real numbers, we are allowing $a = -\infty$ and $b = \infty$.

²If either endpoint a or b is included in I (e.g., $I = [a, b]$), these limits are taken to be one-sided limits when necessary (either $h > 0$ at $x = a$ or $h < 0$ at $x = b$).

³Which is, admittedly, a terrible name.

With these sets at hand, we say that a function f is smooth or infinitely differentiable on I if it belongs to $C^k(I)$ for all k , i.e.,

$$f \in C^\infty(I) := \bigcap_{k=0}^{\infty} C^k(I).$$

We have the following proposition:

Proposition 1.1. *Let f and g be complex-valued functions on I and let α and β be complex numbers.*

1. *If f and g are differentiable on I , then so are $\alpha f + \beta g$, fg and f/g (provided that $g \neq 0$ on I) and we have the formulas*

$$(a) \quad \frac{d}{dx}(\alpha f + \beta g)(x) = \alpha f'(x) + \beta g'(x) \text{ for } x \in I.$$

$$(b) \quad \frac{d}{dx}(fg)(x) = f'(x)g(x) + f(x)g'(x) \text{ for } x \in I.$$

$$(c) \quad \frac{d}{dx}(f/g)(x) = (f'(x)g(x) - f(x)g'(x))/g(x)^2 \text{ for } x \in I.$$

2. *For any $k = 0, 1, 2 \dots \infty$, if $f, g \in C^k(I)$, then $\alpha f + \beta g \in C^k(I)$. In particular, this guarantees that each $C^k(I)$ is a subspace of the vector space of complex-valued functions on I and is therefore a vector space in its own right.*

Though the above proposition might seem obvious (at least the first item) and it does follow from the results of single-variable (real-valued) calculus, you should think about what is meant by all of the algebraic operations above. As a good check, here is a basic exercise:

Exercise 1

Writing $f(x) = u(x) + iv(x)$ and $\alpha = a + ib$. Verify that

$$\frac{d}{dx}(\alpha f)(x) = \alpha f'(x)$$

(whenever f is differentiable) by expanding both sides using complex multiplication and applying the definition of the derivative of a complex-valued function given above.

Exercise 2: The complex exponential function

One of the most important complex-valued functions is the function $x \mapsto e^{ix}$ defined by

$$e^{ix} = \cos(x) + i \sin(x)$$

for $x \in I = \mathbb{R}$.

1. What is e^{i0} ? What is $e^{i\pi}$? Note: If you haven't seen this before, now is a good time to Google "most beautiful equation in Math".
2. Using trigonometric identities (and the algebra of complex numbers), show that

$$e^{i(x+y)} = e^{ix} e^{iy}$$

for any $x, y \in \mathbb{R}$.

3. Upon recalling that, for a non-zero complex number $z = a + ib$, $w = 1/z$ is the unique complex number for which $wz = zw = 1 + 0i$, show that

$$e^{-ix} = \frac{1}{e^{ix}}$$

for each $x \in \mathbb{R}$ (note, by e^{-ix} , I mean $e^{i(-x)}$).

4. Using only the definition above and your knowledge of how complex-numbers are multiplied, show that e^{ix} is differentiable on \mathbb{R} and

$$\frac{d}{dx}e^{ix} = ie^{ix}$$

and conclude that $e^{ix} \in C^1(\mathbb{R})$.

5. Using mathematical induction, show that e^{ix} is k -times differentiable on \mathbb{R} with

$$\frac{d^k}{dx^k}e^{ix} = (i)^k e^{ix}$$

for every $k = 0, 1, 2, \dots$. Conclude that $f \in C^k(\mathbb{R})$ for every k and so $x \mapsto e^{ix}$ is smooth.

1.2 Integral Calculus

In single-variable calculus, you learned to integrate functions $u : I \rightarrow \mathbb{R}$ using limits of Riemann sums. There, you learned that continuous functions were integrable over bounded⁴ intervals I and that integration was a linear operation (just like differentiation). In this short subsection, we carry these ideas over to complex-valued functions of a real variable.

For $f = u + iv \in C^0(I)$, we define its integral over I by the complex number

$$\int_I f = \int_I f(x) dx = \int_I u(x) dx + i \int_I v(x) dx$$

where $\int_I u(x) dx$, $\int_I v(x) dx$ are your (existent) Riemann (or Riemann-Darboux) integrals of Calculus 1. In the case that I is unbounded, these should be taken as improper Riemann integrals (and u and v should additionally be assumed to be improperly Riemann integrable).

Proposition 1.2 (The integral is linear). *Given an interval I , let $f, g \in C^0(I)$ and α and β be complex numbers. Then*

$$\int_I (\alpha f + \beta g) = \alpha \int_I f + \beta \int_I g.$$

Exercise 3

As with the analogous proposition for differentiation, the proposition above seems obvious. Still, to get a feel for what it is saying, please prove the following special case (using only results of single-variable calculus): For a continuous function $f = u + iv \in C^0(I)$ and $\alpha = a + ib \in \mathbb{C}$, show that

$$\int_I \alpha f = \alpha \int_I f.$$

(You may assume I is a bounded interval).

Exercise 4: The fundamental theorem of calculus and integration by parts

Fix $I = [a, b]$. For a function $f = u + iv \in C^0(I)$, we say that $F = U + iV \in C^1(I)$ is an antiderivative for f on I if $F' = f$ on I .

1. If F is an antiderivative of f on I , use the single-variable calculus version of the fundamental theorem

⁴For unbounded intervals, we shall use the notion of Improper Riemann Integral.

of calculus to show that

$$\int_I f = F(x)|_a^b = F(b) - F(a).$$

- Use the above FTC for complex-valued function to prove the following complex-valued intergration by parts formula: For any $f, g \in C^1(I)$,

$$\int f'g dx = f(x)g(x)|_a^b - \int fg'.$$

On the vector space $C^0(I)$, we define an inner product (called the L^2 inner product) as follows: For $g, f \in C^0(I)$,

$$\langle g, f \rangle = \int_I g\bar{f} = \int_I g(x)\overline{f(x)} dx$$

where $\overline{f(x)}$ is the complex-conjugate of the function $f(x) = u(x) + iv(x)$ defined by

$$\overline{f(x)} = u(x) - iv(x)$$

The following proposition (whose details I encourage you to check) shows that $\langle \cdot, \cdot \rangle$ is a *bona fide* complex inner product (like the dot product on \mathbb{C}^n).

Proposition 1.3. *The map $f, g \mapsto \langle f, g \rangle$ is a complex inner product on $C^0(I)$ in that it satisfies the following properties:*

- For $f, g, h \in C^0(I)$,

$$\langle f + h, g \rangle = \langle f, g \rangle + \langle h, g \rangle.$$

and

$$\langle f, g + h \rangle = \langle f, g \rangle + \langle f, h \rangle.$$

- For $\alpha \in \mathbb{C}$ and $f, g \in C^0(I)$,

$$\langle \alpha f, g \rangle = \alpha \langle f, g \rangle$$

and

$$\langle f, \alpha g \rangle = \bar{\alpha} \langle f, g \rangle.$$

- For any $f, g \in C^0(I)$,

$$\langle f, g \rangle = \overline{\langle g, f \rangle}.$$

- For any $f \in C^0$,

$$\langle f, f \rangle \geq 0.$$

This is zero if and only if f is the zero function.

2 Sturm-Liouville Theory

Let p and q be real-valued functions on an interval $I = [a, b]$ and assume, additionally that $p \in C^1(I)$, and $q \in C^0(I)$. For a function $y \in C^2(I)$, define $L[y]$ by

$$L[y](x) = -\frac{d}{dx} (p(x)y'(x)) + q(x)y(x)$$

for $x \in I$. Our goal in this section is to solve the so-called Sturm-Liouville problem: Given $r \in C^0(I)$ for which $r(x) > 0$ on I and constants C_a, C'_a, C_b , and C'_b , determine (the existence of) constants λ for which

$$\begin{cases} L[y] = \lambda r(x)y(x) & x \in I \\ C_a y(a) + C'_a y'(a) = 0 \\ C_b y(b) + C'_b y'(b) = 0, \end{cases} \tag{1}$$

admits a non-zero solution $y = y_\lambda \in C^2(I)$. Further, we aim to determine the nature of these constants and their associated solutions $y = y_\lambda$.

Exercise 5

As defined above, show that L is a linear operator/transformation from $C^2(I)$ to $C^0(I)$. To do this, you'll need to verify that $L[y] \in C^0(I)$ whenever $y \in C^2(I)$ and that the transformation L is linear, i.e., $L[\alpha_1 y_1 + \alpha_2 y_2] = \alpha_1 L[y_1] + \alpha_2 L[y_2]$ for $y_1, y_2 \in C^2(I)$ and $\alpha_1, \alpha_2 \in \mathbb{C}$.

For simplicity, we shall focus on the version of the Sturm-Liouville problem with $r \equiv 1$ which, in this case, we can phrase as an eigenvalue problem. We first need a vector space over which to phrase it. To this end, let

$$\mathcal{D}_L = \{y \in C^2(I) : C_a y(a) + C'_a y'(a) = 0 \text{ and } C_b y(b) + C'_b y'(b) = 0\}.$$

Exercise 6

Show that \mathcal{D}_L is a subspace of $C^2(I)$.

Remark 2.1. It is a very common practice to phrase a boundary-value problem in ODE or PDE in terms of linear algebra (or, technically, functional analysis, linear algebra's bigger sibling). In this case, the boundary conditions are built in to the underlying vector space (which is the domain of the linear operator used to pose the problem). This is a big idea that one gets a lot of mileage out of.

Because \mathcal{D}_L is a subspace of $C^2(I)$, it is a vector space in its own right. In this perspective, solving the Sturm-Liouville Problem (1) (with $r \equiv 1$), is equivalent to finding an eigenvalue λ for the operator L defined on \mathcal{D}_L , i.e., find a complex number λ and a non-zero member/function $y = y_\lambda \in \mathcal{D}_L$ for which

$$L[y](x) = \lambda y(x)$$

for $x \in I$. For this reason, in particular, we refer to such values λ as eigenvalues for (1) and corresponding solutions, y_λ , eigenvectors. To help understand the nature of these eigenvalues/vectors, it's helpful to use some ideas from analysis.

Proposition 2.2 (Lagrange's identity). *For any f and g in $C^2(I)$, we have*

$$\langle L[f], g \rangle - \langle f, L[g] \rangle = \left(p(x) \left(f(x) \overline{g'(x)} - f'(x) \overline{g(x)} \right) \right) \Big|_a^b$$

Proof. Using integration by parts and the linearity of the integral, we see that

$$\begin{aligned} \langle L[f], g \rangle &= \int_I - (p(x) f'(x))' \overline{g(x)} dx + \int_I q(x) f(x) \overline{g(x)} dx \\ &= -p(x) f'(x) \overline{g(x)} \Big|_a^b + \int_I p(x) f'(x) \overline{g(x)}' dx + \int_I q(x) f(x) \overline{g(x)} dx \\ &= -p(x) f'(x) \overline{g(x)} \Big|_a^b + \int_I f'(x) \overline{p(x) g'(x)} dx + \int_I f(x) \overline{q(x) g(x)} dx \end{aligned}$$

where we have used the fact that $p(x)$ and $q(x)$ are real-valued (so that $p(x) \overline{g(x)} = \overline{p(x) g(x)}$ and $q(x) \overline{g(x)} = \overline{q(x) g(x)}$) and also that complex conjugation commutes with differentiation, i.e., $(f)' = f'$. Another application of integration by parts shows that

$$\int_I f'(x) \overline{p(x) g'(x)} dx = f(x) \overline{p(x) g'(x)} \Big|_a^b - \int_I f(x) \overline{(p(x) g'(x))}' dx$$

Putting the two preceding equations together, we obtain

$$\begin{aligned}
 \langle L[f], g \rangle &= -p(x)f'(x)\overline{g(x)} \Big|_a^b + f(x)\overline{p(x)g'(x)} \Big|_a^b - \int_I f'(x)\overline{(p(x)g'(x))'} dx + \int_I f(x)\overline{q(x)g(x)} dx \\
 &= \left(p(x)f(x)\overline{g'(x)} \right) \Big|_a^b - \left(p(x)f'(x)\overline{g(x)} \right) \Big|_a^b + \int_I f(x)\overline{(-p(x)g'(x))' + q(x)g(x)} dx \\
 &= \left(p(x) \left(f(x)\overline{g'(x)} - f'(x)\overline{g(x)} \right) \right) \Big|_a^b + \int_I f(x)\overline{L[g](x)} dx \\
 &= \left(p(x) \left(f(x)\overline{g'(x)} - f'(x)\overline{g(x)} \right) \right) \Big|_a^b + \langle f, L[g] \rangle
 \end{aligned}$$

and from this the proposition follows immediately. □

Corollary 2.3. For $y_1, y_2 \in \mathcal{D}_L$,

$$\langle L[y_1], y_2 \rangle = \langle y_1, L[y_2] \rangle.$$

This is to say that L is a Hermitian operator on the vector space \mathcal{D}_L .

Exercise 7

Prove the corollary above. In view of Lagrange's identity, it suffices to verify that

$$\left(y_1(x)\overline{y_2'(x)} - y_1'(x)\overline{y_2(x)} \right) \Big|_a^b = 0$$

whenever y_1 and y_2 satisfy the boundary conditions:

$$C_a y(a) + C'_a y'(a) = 0 \quad \text{and} \quad C_b y(b) + C'_b y'(b) = 0$$

where C_a, C'_a, C_b , and C'_b are real numbers (you'll need the fact that they are real).

The following proposition shows that our worrying about complex numbers is mostly unnecessary.

Proposition 2.4. If λ is an eigenvalue of (1), then λ is real and it has a real-valued eigenfunction $y_\lambda \in \mathcal{D}_L$.

Proof. We assume that an eigenfunction $f = f_\lambda \in \mathcal{D}_L$ exists and write $f = u + iv$. In view of Lagrange's identity, observe that

$$\lambda \langle f, f \rangle = \langle \lambda f, f \rangle = \langle L[f], f \rangle = \langle f, L[f] \rangle = \langle f, \lambda f \rangle = \overline{\lambda} \langle f, f \rangle$$

and therefore

$$(\lambda - \overline{\lambda}) \langle f, f \rangle = 0.$$

Upon noting that

$$\langle f, f \rangle = \int_I |f(x)|^2 dx = \int_I u(x)^2 + v(x)^2 dx > 0$$

because $f(x)\overline{f(x)} = |f(x)|^2 = u(x)^2 + v(x)^2$ is continuous and non-negative on I , we conclude that

$$\lambda = \overline{\lambda}.$$

This can only happen when λ is a real number (You should check it!). It follows that the equation

$$L[u + iv] = L[f] = \lambda f = \lambda(u + iv)$$

can be split up into its real and imaginary parts and we obtain

$$L[u] = \lambda u \quad \text{and} \quad L[v] = \lambda v$$

(mostly because L has no parts that introduce a phase – it just operates on the real and imaginary parts separately since p and q are real). Since $f = f_\lambda$ is not zero (by its definition as an eigenfunction), u and v cannot both be zero (identically). If $u \neq 0$, put $y_\lambda = u$. Otherwise, set $y_\lambda = v$. By this argument, we have shown that λ is real and has a real-valued eigenfunction $y_\lambda \in C^2(I)$. Finally, we note that the boundary conditions for f can also be split into real and imaginary parts (just as we did with $L[f] = \lambda f$) since the constants $C_a, C'_a, C_b,$ and C'_b are real. From this it follows that $y_\lambda \in \mathcal{D}_L$ and the proof is complete. \square

In view of the preceding proposition, all eigenfunctions for L (corresponding to necessarily real-valued eigenvalues) will be taken to be real-valued.

Proposition 2.5. *Suppose that λ_1 and λ_2 are eigenvalues for (1) with corresponding eigenfunctions $y_1, y_2 \in \mathcal{D}_L$. If $\lambda_1 \neq \lambda_2$, then y_1 and y_2 are orthogonal with respect to the inner product $\langle \cdot, \cdot \rangle$, i.e.,*

$$\langle y_1, y_2 \rangle = 0$$

whenever $\lambda_1 \neq \lambda_2$.

Proof. By assumption, we have that $L[y_1] = \lambda_1 y_1$ and $L[y_2] = \lambda_2 y_2$, then by the preceding corollary, we have

$$\langle \lambda_1 y_1, y_2 \rangle = \langle L[y_1], y_2 \rangle = \langle y_1, L[y_2] \rangle = \langle y_1, \lambda_2 y_2 \rangle = \overline{\lambda_2} \langle y_1, y_2 \rangle$$

and, using the fact that λ_2 is real (i.e., $\lambda_2 = \overline{\lambda_2}$), we rewrite this as

$$(\lambda_1 - \lambda_2) \langle y_1, y_2 \rangle.$$

Thus, if $\lambda_1 \neq \lambda_2$, we must have $\langle y_1, y_2 \rangle = 0$. \square

Theorem 2.6. *The eigenvalues of the Sturm-Liouville problem (1) are all simple, that is, to each eigenvalue λ , its eigenspace*

$$Eig(\lambda) = \{y \in \mathcal{D}_L : L[y] = \lambda y\}$$

is one-dimensional. Further, there exists an infinite sequence of eigenvalues $\{\lambda_n\} \subseteq \mathbb{R}$ which can be put in increasing order, i.e., $\lambda_1 < \lambda_2 < \lambda_3 < \dots$. Furthermore,

$$\lim_{n \rightarrow \infty} \lambda_n = \infty.$$

I'm not sure how much of this I'm going to prove. But, this might be a nice project for someone interested in the analysis aspects of the course. There are a number of ways to prove this theorem; one is using the Riesz theory of compact self-adjoint operators.

3 Life on the circle

Our analysis thus far has focused on complex-valued functions defined on intervals of the form $I = [a, b]$. In our study of Fourier series, we will study complex-valued functions on the “circle” or on the torus. As we see below, this is just a fancy way of talking about complex-valued and 2π -periodic functions – such objects have several equivalent descriptions. As a first introduction, consider the unit circle

$$S^1 = \left\{ z = a + ib \in \mathbb{C} : |z| = \sqrt{a^2 + b^2} = 1 \right\}$$

in the complex plane \mathbb{C} . As a direct consequence of Exercise B (See the Appendix), S^1 is given by

$$S^1 = \{e^{i\theta} : \theta \in \mathbb{R}\}.$$

In fact, the exercise states that

$$S^1 = \{e^{i\theta} : \theta \in (-\pi, \pi]\}$$

and this description is one to one in the sense that the function $\theta \mapsto e^{i\theta}$ bijectively maps $(-\pi, \pi]$ onto the unit circle S^1 . The following proposition gives an equivalence between 2π -periodic functions, functions on the interval $(-\pi, \pi]$ and functions on the circle.

Proposition 3.1. *There is a one-to-one-to-one correspondence between the complex-valued functions on S^1 , the complex-valued functions on $(-\pi, \pi]$ and the complex-valued and 2π -periodic functions on \mathbb{R} . This correspondence is given by*

$$\tilde{f}(\theta) = F(e^{i\theta}) = F(z) = F(e^{ix}) = f(x)$$

where $z = e^{i\theta} = e^{ix}$ for function $F : S^1 \rightarrow \mathbb{C}$, $\tilde{f} : (-\pi, \pi] \rightarrow \mathbb{C}$ and $f : \mathbb{R} \rightarrow \mathbb{C}$, the latter of which is 2π -periodic.

As the proof of the proposition is not terribly illuminating, I'll omit it. Figure 1 illustrates this correspondence and the main idea of the proposition. In view of proposition, we will focus our attention on complex-valued and 2π -periodic functions which we will usually denote by f . These functions will often be said to be functions on the torus \mathbb{T} , however, this is a slight abuse of language and notation, both of which are justified by the proposition. Precisely, the torus $\mathbb{T} = \mathbb{R}/(2\pi\mathbb{Z})$ which can be recognized as the quotient of the additive groups \mathbb{R} and $2\pi\mathbb{Z}$. In fact, one can also recognize \mathbb{T} by the interval $(-\pi, \pi]$. This association is given by the fact that there is a one-to-one correspondence between the interval $(-\pi, \pi]$ and $\mathbb{R}/(2\pi\mathbb{Z})$. You won't however need to worry about this construction nor what a quotient group is. You can think of this reference as merely cultural (where I mean the culture of harmonic analysis in the lens of topological group theory). The essential thing you'll need to understand, and to work with, is the notion of 2π -periodic functions. Just know that there is a lot of important group theory going on in the background.

3.1 Integration on \mathbb{T} and some important spaces of functions on \mathbb{T}

Okay, now let's talk about integration on \mathbb{T} , that is, the integration of 2π periodic functions.

Definition 3.2. *Given a function $f : \mathbb{R} \rightarrow \mathbb{C}$, we say that f is Riemann-integrable on the torus \mathbb{T} (or integrable on the circle) if f is 2π periodic and Riemann-integrable on the interval $[-\pi, \pi]$. In this case we write $f \in R(\mathbb{T})$ and define*

$$\int_{\mathbb{T}} f = \int_{[-\pi, \pi]} f(x) dx;$$

this is called the integral of f over \mathbb{T} . We will also write

$$\int_{\mathbb{T}} f(x) dx = \int_{\mathbb{T}} f.$$

As the integral of f on \mathbb{T} is defined in terms of the Riemann integral on the interval $[-\pi, \pi]$, it's easy to see that the theory on the Riemann-Darboux integral (see the Appendix, and all related tests, theorems, etc) apply to this integral too, with only minor (and obvious) changes in notation. Instead of restating every result in this new notation, we will simply refer to the original results and ask the reader to make appropriate changes to notation and context.

Remark 3.3 (A word of caution). It is common for authors to define the integral of a function f over \mathbb{T} instead by the number

$$\frac{1}{2\pi} \int_{[-\pi, \pi]} f(x) dx.$$

This convention (which we do not use) has some advantages in Fourier analysis as one doesn't have to carry around 2π everywhere and so formulas/statements become simpler. Though I do like this convention, it is slightly less transparent and so I've decided to avoid it. In any case, in your readings of other texts/notes, you should watch out as it is often not clear which convention is being used.

One result that is special for the integral of a function f over \mathbb{T} , is that it can be computed by integrating f over any interval of length 2π , not just the interval $[-\pi, \pi]$. This fact, captured by the following proposition, relies essentially on our requirement that f is 2π -periodic.




Figure 1: An illustration of the correspondence between functions on the circle, functions on $(-\pi, \pi]$ and 2π -periodic functions

Proposition 3.4. *Let $f : \mathbb{R} \rightarrow \mathbb{C}$ be 2π -periodic. Then $f \in R(\mathbb{T})$ if and only if, $f \in R(I)$ for some interval I of*

length 2π , i.e., $I = [a, a + 2\pi]$ for some $a \in \mathbb{R}$. For any such f , we have

$$\int_{\mathbb{T}} f(x) dx = \int_I f(x) dx.$$

In other words, integrating f over any interval of length 2π gives the same number.

Exercise 8

Suppose that $f : \mathbb{R} \rightarrow \mathbb{C}$ is 2π -periodic. With Proposition B.10 in mind, do the following.

1. Prove the following statement. If $f \in R([a, b])$ then, for any $n \in \mathbb{Z}$, $f \in R([a + 2\pi n, b + 2\pi n])$ and

$$\int_a^b f = \int_{a+2\pi n}^{b+2\pi n} f.$$

2. Prove the following statement. If $f \in R([-\pi, \pi])$, then, for any $x_0 \in \mathbb{R}$, the function $x \mapsto f(x + x_0)$ is Riemann integrable on $[-\pi, \pi]$ and

$$\int_{-\pi}^{\pi} f(x) dx = \int_{-\pi}^{\pi} f(x + x_0) dx.$$

3. Prove Proposition 3.4.

In view of Definition 3.2, $R(\mathbb{T})$ is the set of Riemann integrable functions on \mathbb{T} . Structurally, this set can be recognized a vector space (over \mathbb{C}) of complex-valued functions under the usual notion of function addition. Also, $f \mapsto \int_{\mathbb{T}} f$ is an important linear map from this vector space into the complex plane. Let's introduce some other important spaces of functions, all of which turn out to be subspaces of $R(\mathbb{T})$.

Definition 3.5. Let $f : \mathbb{R} \rightarrow \mathbb{C}$ be 2π periodic.

1. We say that f is continuous on \mathbb{T} if f is continuous on \mathbb{R} . Here we write that $f \in C(\mathbb{T})$ or $f \in C^0(\mathbb{T})$.
2. We say that f is once continuously differentiable on \mathbb{T} if f is differentiable on \mathbb{R} with continuous derivative $f' \in C^0(\mathbb{T})$. In this case we write $f \in C^1(\mathbb{T})$.
3. More generally, we say that f is n -times continuously differentiable on \mathbb{T} if $f \in C^{n-1}(\mathbb{T})$ and the $(n-1)$ th derivative of f , $f^{(n-1)}$, is differentiable and its derivative $f^{(n)} := (f^{(n-1)})' \in C(\mathbb{T})$. In this case, we write $f \in C^n(\mathbb{T})$.
4. We say that f is smooth on \mathbb{T} if $f \in C^n(\mathbb{T})$ for all n . In this case we write $f \in C^\infty(\mathbb{T})$.

Exercise 9

To clarify the preceding definition, this exercise asks you to work out some details. First, let's fix something that the definition sweeps under the carpet, so to speak.

1. Given a 2π -periodic function f , it makes sense to ask its derivative f' exists and is continuous which is what we usually mean by continuously differentiable. The definition above requires, however, that $f' \in C^0(\mathbb{T})$ and so it requires the additional condition that f' is also a 2π periodic function. To clear this up, prove the following statement:

If $f : \mathbb{R} \rightarrow \mathbb{C}$ is 2π -periodic and differentiable with derivative $f' : \mathbb{R} \rightarrow \mathbb{C}$. Then f' is also 2π -periodic.

Given a 2π -periodic function $f : \mathbb{R} \rightarrow \mathbb{C}$, for $f \in C^1(\mathbb{T})$ the definition above requires that f is differentiable on all of the real line \mathbb{R} and, moreover, that f' is continuous. For the next item, you will prove that this requirement is unnecessarily strong.

2. Prove that $f \in C^1(\mathbb{T})$ if and only if f is continuously differentiable (meaning it is differentiable and has continuous derivative) on some interval $I = (a, b)$ where $b - a > 2\pi$.
3. Show that the equivalent condition of the above statement cannot be relaxed. That is, find a function $f : \mathbb{R} \rightarrow \mathbb{C}$ which is 2π -periodic and for which f is continuous differentiable on some interval $I = (a, b)$ for which $b - a = 2\pi$, yet $f \notin C^1(\mathbb{T})$.

As continuous functions are integrable and differentiable functions are continuous (and so on and so forth), we obtain

$$C^\infty(\mathbb{T}) \subseteq \dots \subseteq C^n(\mathbb{T}) \subseteq \dots \subseteq C^1(\mathbb{T}) \subseteq C^0(\mathbb{T}) \subseteq R(\mathbb{T})$$

where each of these containments is proper. All of the above sets are, in fact, vector spaces over \mathbb{C} under the usual notion of function addition as we previously discussed and, further, all of above relations remain true when one replaces “ \subseteq ” with “ \leq ” where $V \leq W$ means V is a subspace of W .

3.2 Pointwise and Uniform convergence of functions on \mathbb{T}

Throughout the next **several** subsections, we explore several notions of convergence of Fourier series for functions on \mathbb{T} . To this end, this subsection is dedicated to stating (and restating) several notions and results pertaining to the convergence of functions and series of functions on \mathbb{T} . The functions (and sequences of functions) we study will be taken to be, at worst, Riemann-integrable on \mathbb{T} .

Consider a sequence of functions $\{f_n\}_{n=1}^\infty \subseteq R(\mathbb{T})$ and another function $f \in R(\mathbb{T})$. Given a set $E \subseteq \mathbb{R}$, we say that $\{f_n\}$ *converges to f on E* if, for each $x \in E$, the sequence of complex numbers $\{f_n(x)\}$ converges to $f(x)$, i.e., $\lim_{n \rightarrow \infty} f_n(x) = f(x)$. As all functions in sight are 2π -periodic, $\lim_{n \rightarrow \infty} f_n(x) = f(x)$ if and only if $\lim_{n \rightarrow \infty} f_n(x + 2\pi k) = f(x + 2\pi k)$ for each $k \in \mathbb{Z}$. In other words, $\{f_n\}$ converges to f on E if and only if $\{f_n\}$ converges on the set

$$\bigcup_{k \in \mathbb{Z}} (E + 2\pi k) = \{x \in \mathbb{R} : x = y + 2\pi k \text{ for some } y \in E \text{ and } k \in \mathbb{Z}\}.$$

In the special case that $\{f_n\}$ converges to f on \mathbb{R} or, equivalently, on any interval of length at least 2π , we say that $\{f_n\}$ converges pointwise to f on \mathbb{T} . In other words, $\{f_n\}$ converges pointwise to f on \mathbb{T} if, for each $x \in \mathbb{R}$ and $\epsilon > 0$, there exists a natural number N for which

$$|f_n(x) - f(x)| < \epsilon$$

whenever $n \geq N$. Just as we did on an interval I ,

Definition 3.6. *Let $\{f_n\}$ be a sequence of functions in $R(\mathbb{T})$ and let $f \in R(\mathbb{T})$. We say that $\{f_n\}$ converges uniformly to f on \mathbb{T} if, for each $\epsilon > 0$, there exists a natural number N for which*

$$|f_n(x) - f(x)| < \epsilon$$

for all $n \geq N$ and $x \in \mathbb{R}$.

Of course, the above notion of uniform convergence was the same as it was for uniform convergence of functions on \mathbb{R} . The difference here is that the functions of interest are 2π periodic. In fact, it's easy to see that $\{f_n\}$ converges uniformly to f on \mathbb{T} if and only if $\{f_n\}$ converges uniformly to f on any interval I of length at least 2π . As we saw before, uniform convergence on \mathbb{T} is also captured by the sup norm. Let's define it in this context.

Definition 3.7. Given any $f \in R(\mathbb{T})$ (which is necessarily bounded), define

$$\|f\|_\infty = \sup_{\mathbb{T}} |f(x)|$$

where $\sup_{\mathbb{T}} |f(x)| := \sup\{|f(x)| : x \in I\}$ given any interval $I \subseteq \mathbb{R}$ of length at least 2π .

In this language and by virtue of our previous results, we have the following proposition.

Proposition 3.8. Let $\{f_n\} \in R(\mathbb{T})$ and let $f \in R(\mathbb{T})$. Then $\{f_n\}$ converges uniformly to f on \mathbb{T} if and only if

$$\lim_{n \rightarrow \infty} \|f_n - f\|_\infty = 0.$$

If either of the equivalent conditions of the above proposition is met, $\{f_n\}$ is also said to converge to f with respect to the $L^\infty(\mathbb{T})$ norm – a notion we will discuss more fully in the next subsection. In the context of the torus \mathbb{T} , we restate Theorem C.4:

Theorem 3.9. Let $\{f_n\}$ be a sequence of functions in $R(\mathbb{T})$. Then $\{f_n\}$ converges uniformly on \mathbb{T} if and only if the following condition is satisfied:

(UC) For all $\epsilon > 0$, there exists a natural number N such that

$$|f_n(x) - f_m(x)| < \epsilon \quad \text{whenever } x \in \mathbb{R} \text{ and } n, m \geq N.$$

Let's also recapture Theorems C.7 and C.8 in the context of the torus \mathbb{T} .

Theorem 3.10. Let $\{f_n\}$ be a sequence of functions in $R(\mathbb{T})$ and suppose that $\{f_n\}$ converges uniformly on \mathbb{R} to a function $f : \mathbb{R} \rightarrow \mathbb{C}$. Then the following properties hold:

1. The limit f is Riemann integrable on \mathbb{T} , i.e., $f \in R(\mathbb{T})$, and the sequence $\{f_n\}$ converges uniformly to f on \mathbb{T} .
2. If $\{f_n\} \subseteq C(\mathbb{T})$, then $f \in C(\mathbb{T})$.
3. We have

$$\lim_{n \rightarrow \infty} \int_{\mathbb{T}} f_n = \int_{\mathbb{T}} \lim_{n \rightarrow \infty} f_n = \int_{\mathbb{T}} f.$$

Let's now talk about series on \mathbb{T} . Given a sequence $\{f_n\} \in R(\mathbb{T})$, we investigate the corresponding series $\sum_n f_n$. For each natural number N , we define the N th partial sum S_N of the series $\sum_n f_n$ by

$$S_N(x) = \sum_{n=1}^N f_n(x)$$

for $x \in \mathbb{R}$. It is clear that $\{S_N\}$ is a sequence of Riemann-integrable functions on \mathbb{T} and we can ask whether or not it has a limit. If, for $x \in \mathbb{R}$,

$$\lim_{N \rightarrow \infty} S_N(x) = \lim_{N \rightarrow \infty} \sum_{n=1}^N f_n(x)$$

exists, we say that the series $\sum_n f_n$ converges at x and define the sum of the series to be the number

$$\sum_{n=1}^{\infty} f_n(x) = \lim_{N \rightarrow \infty} S_N(x) = \lim_{N \rightarrow \infty} \sum_{n=1}^N f_n(x).$$

If the sequence of partial sums S_N converges at each $x \in \mathbb{R}$, we say that the series $\sum_{n=1}^{\infty} f_n$ converges pointwise on \mathbb{T} and write

$$\sum_{n=1}^{\infty} f_n(x) = \lim_{N \rightarrow \infty} S_N(x)$$

for each $x \in \mathbb{R}$. Further, we say that the series $\sum_{n=1}^{\infty} f_n$ converges uniformly on \mathbb{T} if the sequence $\{S_N\}$ of partial sums converges uniformly on \mathbb{T} . Rephrasing our work from the previous section, we can state the Weierstrass M -test in the context of \mathbb{T} .

Theorem 3.11. *Let $\{f_n\}$ be a sequence of functions in $R(\mathbb{T})$ and, for each $n \in \mathbb{N}$, let $M_n = \|f_n\|_\infty$. If the numerical series $\sum_{n=1}^\infty M_n$ converges, then the following statements hold:*

1. *The series $\sum_{n=1}^\infty f_n$ converges uniformly on \mathbb{T} .*

2. *We have*

$$\int_{\mathbb{T}} \sum_{n=1}^\infty f_n(x) dx = \sum_{n=1}^\infty \int_{\mathbb{T}} f_n(x) dx.$$

3. *If, additionally, $\{f_n\} \subseteq C(\mathbb{T})$, the sum of the series*

$$\sum_{n=1}^\infty f_n(x) = \lim_{N \rightarrow \infty} \sum_{n=1}^N f_n(x)$$

is also a member of $C(\mathbb{T})$.

3.3 The Lebesgue norms on \mathbb{T} and the $L^2(\mathbb{T})$ inner product

We begin this section by introducing the Lebesgue norms on the torus \mathbb{T} . These norms, denoted by $\|\cdot\|_p$ for $1 \leq p \leq \infty$, give a notion of “size” to each function in $R(\mathbb{T})$ and, in fact, we’ve already seen some beautiful properties of the $\|\cdot\|_\infty$ norm studied in the preceding section. As we did on subintervals of the real line, we will pay special attention to the $p = 2$ case as it is captured by the rich structure of the so-called L^2 inner product.

Warning: The notation in this subsection will parallel that of Subsection C.3; however, the definitions will all differ by a factor of $(2\pi)^{1/p}$. These changes are only to simplify notation and won’t affect anything essentially. In any case, watch out for the 2π !

Definition 3.12. 1. *For $1 \leq p < \infty$, we define the $L^p = L^p(\mathbb{T})$ norm of a function $f \in R(\mathbb{T})$ by*

$$\|f\|_p = \left(\frac{1}{2\pi} \int_{\mathbb{T}} |f(x)|^p dx \right)^{1/p}.$$

2. *If $p = \infty$, we define the $L^p = L^\infty = L^\infty(\mathbb{T})$ of $f \in R(\mathbb{T})$ by*

$$\|f\|_\infty = \sup_{\mathbb{T}} |f(x)|$$

where $\sup_{\mathbb{T}} |f(x)| := \sup\{|f(x)| : x \in I\}$ given any subinterval $I \subseteq \mathbb{R}$ of length at least 2π , a notion which makes sense precisely because f is 2π -periodic.

3. *Let $1 \leq p \leq \infty$. Given a sequence of function $\{f_n\} \subseteq R(\mathbb{T})$ and another function $f \in R(\mathbb{T})$, we say that $\{f_n\}$ converges to f with respect to the $L^p = L^p(\mathbb{T})$ norm if*

$$\lim_{n \rightarrow \infty} \|f_n - f\|_p = 0.$$

Tracking carefully the appearance of 2π in the above definitions, an appeal to the results of Subsection C.3 yields the following results about the L^p norm on \mathbb{T} .

Proposition 3.13. *For any $f, g \in R(\mathbb{T})$ and $\alpha \in \mathbb{C}$, we have*

1.

$$\|f\|_p \geq 0$$

2.

$$\|\alpha f\|_p = |\alpha| \|f\|_p$$

3.

$$\|f + g\|_p \leq \|f\|_p + \|g\|_p$$

Proposition 3.14 (Hölder's inequality). *Let $1 \leq p \leq q \leq \infty$ be conjugate exponents, i.e.,*

$$\frac{1}{p} + \frac{1}{q} = 1.$$

For any $f, g \in R(\mathbb{T})$,

$$\left| \frac{1}{2\pi} \int_{\mathbb{T}} f(x)g(x) dx \right| \leq \|f\|_p \|g\|_q.$$

Proposition 3.15. *Given any $1 \leq r \leq s \leq \infty$, if a sequence $\{f_n\} \subseteq R(\mathbb{T})$ converges to $f \in R(\mathbb{T})$ with respect to the $L^s(\mathbb{T})$ norm, then $\{f_n\}$ converges to f with respect to the $L^r(\mathbb{T})$ norm. In particular, if $\{f_n\}$ converges to f uniformly, i.e.,*

$$\lim_{n \rightarrow \infty} \|f_n - f\|_{\infty} = 0$$

then, for every $1 \leq p \leq \infty$, $\{f_n\}$ converges to f with respect to the $L^p(\mathbb{T})$ norm, i.e.,

$$\lim_{n \rightarrow \infty} \|f_n - f\|_p = 0.$$

We now turn our focus to the case in which $p = 2$. Here, the norm $\|\cdot\|_2$ is captured by the following inner product.

Definition 3.16. *Given $f, g \in R(\mathbb{T})$, we defined the $L^2 = L^2(\mathbb{T})$ inner product of f and g to be the complex number*

$$\langle f, g \rangle = \frac{1}{2\pi} \int_{\mathbb{T}} f(x) \overline{g(x)} dx$$

Following directly from the definition, we see that the $L^2(\mathbb{T})$ inner product characterizes the $L^2(\mathbb{T})$ norm. That is, for each $f \in R(\mathbb{T})$,

$$\|f\|_2 = \sqrt{\langle f, f \rangle}.$$

Let's summarize some properties of the L^2 inner product.

Proposition 3.17. *Given any $f, g, h \in R(\mathbb{T})$ and $\alpha \in \mathbb{C}$, we have:*

1.

$$\langle f, g \rangle = \overline{\langle g, f \rangle}$$

2.

$$\langle \alpha f, g \rangle = \alpha \langle f, g \rangle$$

3.

$$\langle f, \alpha g \rangle = \overline{\alpha} \langle f, g \rangle$$

4.

$$\langle f + g, h \rangle = \langle f, h \rangle + \langle g, h \rangle$$

5.

$$\langle f, g + h \rangle = \langle f, g \rangle + \langle f, h \rangle$$

Two functions $f, g \in R(\mathbb{T})$ are said to be *orthogonal* (with respect to the L^2 inner product) if $\langle f, g \rangle = 0$. In this case we write $f \perp g$. Using this language, we state three more crucial properties of the $L^2(\mathbb{T})$ inner product and norm.

Proposition 3.18. *Given $f, g \in R(\mathbb{T})$, we have the following statements:*

1. If $f \perp g$, then

$$\|f + g\|_2^2 = \|f\|_2^2 + \|g\|_2^2.$$

2.

$$\left| \frac{1}{2\pi} \int_{\mathbb{T}} f(x)g(x) dx \right| \leq \|f\|_2 \|g\|_2$$

3.

$$\|f + g\|_2 \leq \|f\|_2 + \|g\|_2$$

Proof. We prove only the first property as the others follow from our results of Subsection C.3. Fix $f, g \in R(\mathbb{T})$ and suppose that $f \perp g$. Then, given the properties of the inner product summarized in the preceding proposition,

$$\begin{aligned} \|f + g\|_2^2 &= \langle f + g, f + g \rangle \\ &= \langle f, f + g \rangle + \langle g, f + g \rangle \\ &= \langle f, f \rangle + \langle f, g \rangle + \langle g, f \rangle + \langle g, g \rangle \\ &= \|f\|_2^2 + \langle f, g \rangle + \overline{\langle f, g \rangle} + \|g\|_2^2 \\ &= \|f\|_2^2 + 0 + 0 + \|g\|_2^2 \\ &= \|f\|_2^2 + \|g\|_2^2. \end{aligned}$$

□

Example 1

For any $n \in \mathbb{Z}$, the function $x \rightarrow e^{inx}$ is $C^\infty(\mathbb{T})$ and therefore Riemann integrable on \mathbb{T} . We have, for each $n, m \in \mathbb{Z}$,

$$\langle e^{inx}, e^{imx} \rangle = \frac{1}{2\pi} \int_{\mathbb{T}} e^{inx} e^{-imx} dx = \begin{cases} 1 & n = m \\ 0 & m \neq n \end{cases}$$

as you showed in Homework 1. For this reason, $e^{inx} \perp e^{imx}$ whenever $n \neq m$. Such a collection $\{e^{inx}\}_{n \in \mathbb{Z}}$ is said to be an *orthonormal system*. The word normal comes from the fact that that $L^2(\mathbb{T})$ norm of each function e^{inx} is one.

3.4 Fourier coefficients and the uniform theory

We now (and finally) begin our study of Fourier series. Our main goal is to approximate a given function $f \in R(\mathbb{T})$ by a sequence of trigonometric polynomials, called Fourier polynomials. A trigonometric polynomial of order N is, by definition, a function of the form

$$P_N(x) = \sum_{n=-N}^N c_n e^{inx}$$

for $x \in \mathbb{R}$ where $\{c_n\}$ are complex numbers. It is clear that each such polynomial P_N is a member of $C^\infty(\mathbb{T})$ and hence a member of $R(\mathbb{T})$. Our goal is then to find a sequence of polynomials P_N which approximate f in some sense (pointwise, uniform, with respect to L^2 , etc.). As we will see shortly, the following definition provides a good start.

Definition 3.19. Given $f \in R(\mathbb{T})$, define

$$\hat{f}(n) = \langle f, e^{inx} \rangle = \frac{1}{2\pi} \int_{\mathbb{T}} f(x) e^{-inx} dx$$

for each $n \in \mathbb{Z}$. For each $N \in \mathbb{N}$, we define the Fourier polynomial of order N by

$$S_N(x) = S_N(f)(x) = \sum_{n=-N}^N \hat{f}(n) e^{inx}$$

for $x \in \mathbb{R}$.

The following theorem states that the Fourier coefficients determine a function at each point of continuity. This was proved earlier in the course; for a proof see Theorem 2.1 of the course textbook. We will recapture the result shortly. **I don't like that this is here. Perhaps instead we should do some examples here to show that the Fourier polynomials tell us something about f ?**

Proposition 3.20. Suppose that, for $f, g \in R(\mathbb{T})$, $\hat{f}(n) = \hat{g}(n)$ for all $n \in \mathbb{Z}$. If f and g are continuous at $x \in \mathbb{R}$, then $f(x) = g(x)$. If, additionally, $f, g \in C(\mathbb{T})$, then $f = g$.

With the above definition in mind, we are interested in whether or not the Fourier polynomials of f approximate f in any sense. The natural thing to do is then to take $N \rightarrow \infty$ and this leads us to the notion of Fourier series.

Definition 3.21. given a function $f \in R(\mathbb{T})$, its Fourier series is the formal expression

$$\sum_{n \in \mathbb{Z}} \hat{f}(n) e^{inx} = \sum_{n=-\infty}^{\infty} \hat{f}(n) e^{inx}$$

is called the Fourier series of f . We write

$$f \sim \sum_{n \in \mathbb{Z}} \hat{f}(n) e^{inx}$$

To investigate the convergence of this series, we must investigate the convergence of the Fourier polynomials S_N which are, by definition, partial sums of the Fourier series for f . That is, we say that the Fourier series for f converges at $x \in \mathbb{R}$ if $\lim_{N \rightarrow \infty} S_N(x)$ exists. In this case,

$$\sum_{n \in \mathbb{Z}} \hat{f}(n) e^{inx} = \lim_{N \rightarrow \infty} \sum_{n=-N}^N \hat{f}(n) e^{inx}$$

is the sum of the series. To refine our discussion above, our main goal is to ask: When and in what sense is

$$f(x) = \sum_{n \in \mathbb{Z}} \hat{f}(n) e^{inx}?$$

In view of the Weierstrass M -test, we can give a sufficient condition for the uniform convergence of Fourier series in terms of the summability of the numerical series $\{\hat{f}(n)\}$.

Theorem 3.22. Let $f \in C(\mathbb{T})$ and let $\{\hat{f}(n)\}_{n \in \mathbb{Z}}$ be the Fourier coefficients of f . If the series $\sum_{n \in \mathbb{Z}} |\hat{f}(n)|$ converges, then the Fourier series of f , $\sum_{n \in \mathbb{Z}} \hat{f}(n) e^{inx}$, converges uniformly to f .

Proof. We observe that

$$|\hat{f}(n) e^{inx}| = |\hat{f}(n)| |e^{inx}| = |\hat{f}(n)|$$

for all $n \in \mathbb{Z}$ and $x \in \mathbb{R}$. In other words, the summands $g_n(x) = \hat{f}(n) e^{inx}$ satisfy $|g_n(x)| \leq M_n = |\hat{f}(n)|$ for all $n \in \mathbb{Z}$ and $x \in \mathbb{R}$. In view of our supposition that the series $\sum |\hat{f}(n)| = \sum M_n$ converges, the Weierstrass M -test

guarantees that the Fourier series converges uniformly to some complex-valued function g on \mathbb{R} . Further, because each summand $\hat{f}(n)e^{inx}$ is continuous, g is continuous. Also, observe that

$$g(x + 2\pi) = \lim_{N \rightarrow \infty} \sum_{n=-N}^N \hat{f}(n)e^{in(x+2\pi)} = \lim_{N \rightarrow \infty} \sum_{n=-N}^N \hat{f}(n)e^{inx} = g(x)$$

for all $x \in \mathbb{R}$ and so g is 2π -periodic. We therefore conclude that $g \in C(\mathbb{T})$. It remains to show that $g = f$. To this end, we compute the Fourier coefficients of g . Observe that, for any $m \in \mathbb{Z}$,

$$g(x)e^{-imx} = e^{-imx} \lim_{N \rightarrow \infty} \sum_{n=-N}^N \hat{f}(n)e^{inx} = \lim_{N \rightarrow \infty} \sum_{n=-N}^N \hat{f}(n)e^{inx}e^{-imx}$$

for all $x \in \mathbb{R}$. In fact, the Weierstrass M -text applied to the summands $n \mapsto \hat{f}(n)e^{inx}e^{-imx}$ shows that the partial sums on the right hand side converge uniformly to $g(x)e^{-imx}$. Since each summand is Riemann integrable (it is continuous) and the series converges uniformly, we may integrate term-by-term. For each $m \in \mathbb{Z}$, we have

$$\hat{g}(m) = \langle g, e^{imx} \rangle = \frac{1}{2\pi} \int_{\mathbb{T}} g(x)e^{-imx} dx = \sum_{n \in \mathbb{Z}} \frac{1}{2\pi} \int_{\mathbb{T}} (\hat{f}(n)e^{inx}e^{-imx}) dx = \sum_{n \in \mathbb{Z}} \hat{f}(n) \langle e^{inx}, e^{imx} \rangle = \hat{f}(m)$$

where we have used the fact that $\langle e^{inx}, e^{imx} \rangle = 1$ when $n = m$ and 0 otherwise. Therefore $\hat{f}(n) = \hat{g}(n)$ for all $n \in \mathbb{Z}$ and, in view of the preceding proposition $f = g$. In other words,

$$f(x) = \sum_{n \in \mathbb{Z}} \hat{f}(n)e^{inx}$$

where the convergence is uniform for $x \in \mathbb{R}$. □

I've always found the result above somewhat unsatisfying, though it is powerful as we will shortly see. The reason I find it unsatisfying is because its hypotheses are stated in terms of the Fourier coefficients of f and so, to apply the theorem, one has to compute the Fourier coefficients of f and then ask if they are absolutely summable. One would like to instead have hypotheses stated in terms of f itself. In any case, the theorem allows us to prove the following result.

Corollary 3.23. *If $f \in C^2(\mathbb{T})$, then the Fourier series for f , $\sum_{n \in \mathbb{Z}} \hat{f}(n)e^{inx}$, converges uniformly to f on \mathbb{R} .*

Proof. Let $f(x) = u(x) + iv(x)$ where u and v are real-valued 2π periodic functions which are, by hypothesis, twice continuously differentiable on \mathbb{R} . We have $f'(x) = u'(x) + iv'(x)$ and $f''(x) = u''(x) + iv''(x)$ for $x \in \mathbb{R}$. I claim that, for all non-zero $n \in \mathbb{Z}$,

$$\hat{f}(n) = \frac{-1}{2\pi n^2} \int_{\mathbb{T}} f''(x)e^{-inx} dx.$$

To see this, we apply the complex-version of integration by parts (which works in view of the FTC you proved in Homework 1). This is the statement: If u and v are complex-valued functions differentiable on the interval $[a, b]$ with derivatives u' and v' ,

$$\int_{[a,b]} u(x)v'(x) dx = u(x)v(x) \Big|_a^b - \int_{[a,b]} u'(x)v(x) dx$$

where $u(x)v(x) \Big|_a^b := u(b)v(b) - u(a)v(a)$. So, upon taking $u = f$ and $v'(x) = e^{-inx}$, we have

$$\begin{aligned} \hat{f}(n) &= \frac{1}{2\pi} \int_{\mathbb{T}} f(x)e^{-inx} dx = \frac{1}{2\pi} \int_{[-\pi,\pi]} f(x)e^{-inx} dx \\ &= \frac{1}{2\pi} \left(f(x) \frac{e^{-inx}}{-in} \Big|_{-\pi}^{\pi} - \int_{[-\pi,\pi]} f'(x) \frac{e^{-inx}}{-in} dx \right) \end{aligned}$$

We note, however, that because f and e^{-inx} are 2π periodic, □

Exercise 10

This exercise concerns the absolute decay of Fourier coefficients. It allows you to investigate conditions under which the Fourier series for a function f converges uniformly to f by Theorem 3.22. Do the following:

1. Use induction (on m) and integration by parts to prove the statement: If $f \in C^m(\mathbb{T})$ for $m \in \mathbb{N}$, then, for any non-zero $n \in \mathbb{Z}$,

$$\hat{f}(n) = (in)^{-m} \widehat{f^{(m)}}(n) = \frac{(in)^{-m}}{2\pi} \int_{\mathbb{T}} f^{(m)}(x) e^{-inx} dx$$

where $f^{(m)}$ means the m th-derivative of f . Conclude directly that the Fourier series for f converges uniformly to f whenever $f \in C^m(\mathbb{T})$ for $m \geq 2$.

Given a function $f \in R(\mathbb{T})$, we say that f is Hölder continuous of order $\alpha > 0$ if $f \in C^m(\mathbb{T})$ for $m = \lceil \alpha \rceil - 1$ and there exists $C_\alpha > 0$ for which

$$|f^{(m)}(x) - f^{(m)}(y)| \leq C|x - y|^{\alpha - m}$$

for all $x, y \in \mathbb{R}$.

2. Find a function $f \in R(\mathbb{T})$ which is Hölder continuous of order $\alpha = 2$.
3. Use the mean value theorem to prove that, for $f \in C^1(\mathbb{T})$, f is Hölder continuous of order $\alpha = 1$.
4. Using your reasoning from the previous item, what can be said about the Hölder continuity of f for $f \in C^2(\mathbb{T})$?
5. Prove that, if $f \in R(\mathbb{T})$ is Hölder continuous of order $\alpha > 0$, then there is a constant C_α for which

$$|\hat{f}(n)| \leq \frac{C_\alpha}{|n|^\alpha}$$

for nonzero $n \in \mathbb{Z}$. **Hint:** First differentiate m times for $m = \lceil \alpha \rceil - 1 < \alpha$. In view of your first item, for $n \neq 0$, the 2π -periodicity of $f^{(m)}$ guarantees that

$$\hat{f}(n) = \frac{(in)^{-m}}{2\pi} \int_{-\pi}^{\pi} f^{(m)}(x) e^{-inx} dx = \frac{(in)^{-m}}{2\pi} \int_{-\pi}^{\pi} f^{(m)}\left(x + \frac{\pi}{n}\right) e^{-in(x+\pi/n)} dx.$$

Upon noting that $e^{-in(x+\pi/n)} = -e^{-inx}$, you can average the above integrals to obtain a nice way to write \hat{f} as an integral over a difference $f^{(m)}(x) - f^{(m)}(x + \pi/n)$. Now use Hölder continuity.

6. Use your result from the item above and Theorem 3.22 to prove that the Fourier series for f converges uniformly to f whenever f is Hölder continuous of order α for any $\alpha > 1$. Hint: Appeal to the summability of p series.

This concludes our investigation of the uniform convergence of Fourier series. We now move on to the theory of pointwise convergence. In contrast to our study of uniform convergence, the main object in our study of pointwise convergence of Fourier series revolves around a careful (analytical) study of the Dirichlet kernel, which we introduce now.

Proposition 3.24. Let $f \in R(\mathbb{T})$ and let $(\hat{f}(n))_{n \in \mathbb{Z}}$ be its Fourier coefficients. For each N , let S_N denote the N th Fourier polynomial of f , i.e.,

$$S_N(x) = \sum_{n=-N}^N \hat{f}(n) e^{inx}.$$

Then, for each $N \in \mathbb{N}$ and $x \in \mathbb{R}$,

$$S_N(x) = \frac{1}{2\pi} \int_{\mathbb{T}} D_N(x-y) f(y) dy$$

where

$$D_N(y) := \begin{cases} \frac{\sin((N+1/2)y)}{\sin y/2} & y \neq 2\pi k, k \in \mathbb{Z} \\ 2N+1 & y = 2\pi k, k \in \mathbb{Z} \end{cases} = \sum_{n=-N}^N e^{iny};$$

this is called the Dirichlet kernel. For each N ,

$$\frac{1}{2\pi} \int_{\mathbb{T}} D_N(y) dy = 1$$

and

$$\frac{1}{2\pi} \int_{[-\pi, 0]} D_N(y) dy = \frac{1}{2\pi} \int_{[0, \pi]} D_N(y) dy = \frac{1}{2}.$$

Proof. By virtue of the linearity of the integral, it is evident that

$$\begin{aligned} S_N(x) &= \sum_{n=-N}^N \left(\frac{1}{2\pi} \int_{\mathbb{T}} f(y) e^{-iny} dy \right) e^{inx} \\ &= \frac{1}{2\pi} \int_{\mathbb{T}} \sum_{n=-N}^N f(y) e^{in(x-y)} dy \end{aligned}$$

for $N \in \mathbb{N}$ and $x \in \mathbb{R}$. We now show that

$$D_N(y) = \sum_{n=-N}^N e^{iny}$$

for $N \in \mathbb{N}$ and $x \in \mathbb{R}$. First, it is clear that, if $y = 2\pi k$ for $k \in \mathbb{Z}$, $D_N(y) = 2N+1 = \sum_{n=-N}^N 1 = \sum_{n=-N}^N e^{iny}$. Thus, we assume without loss of generality that $y \neq 2\pi k$ for $k \in \mathbb{Z}$ and hence $\sin y/2 \neq 0$. Now,

$$\begin{aligned} D_1(y) &= \frac{\sin((1+1/2)y)}{\sin(y/2)} = \frac{\sin y \cos(y/2) + \sin(y/2) \cos y}{\sin(y/2)} \\ &= \frac{\sin((1/2+1/2)y) \cos(y/2)}{\sin(y/2)} + \cos y = \frac{2 \sin(y/2) \cos(y/2) \cos(y/2)}{\sin(y/2)} + \cos y \\ &= 2 \cos^2(y/2) + \cos y = 1 + 2 \cos y \\ &= e^{i \cdot 0 \cdot y} + e^{iy} + e^{iy} = \sum_{n=-1}^1 e^{iny} \end{aligned}$$

where we have used the half-angle identity $2 \cos^2(A/2) = 1 + \cos(A)$. Thus, the desired result is true for $N = 1$.

Let's induct on N . Let's assume that the formula holds for $N \geq 1$, we will show it holds for $N + 1$. We have

$$\begin{aligned}
D_{N+1}(y) &= \frac{\sin((N+1+1/2)y)}{\sin y/2} \\
&= \frac{\sin y \cos((N+1/2)y) + \cos y \sin((N+1/2)y)}{\sin y/2} \\
&= 2 \cos(y/2) \cos((N+1/2)y) + \cos y D_N(y) \\
&= 2 \cos(y/2) (\cos Ny \cos(y/2) - \sin Ny \sin(y/2)) + \cos y D_N(y) \\
&= (1 + \cos y) \cos Ny - \sin Ny \sin y + \cos y D_N(y) \\
&= \cos Ny + (\cos y \cos Ny - \sin Ny \sin y) + \cos y D_N(y) \\
&= \cos Ny + \cos((N+1)y) + \cos y D_N(y) \\
&= \frac{e^{iNy} + e^{-iNy}}{2} + \frac{e^{i(N+1)y} + e^{-i(N+1)y}}{2} + \frac{e^{iy} + e^{-iy}}{2} D_N(y) \\
&= \frac{e^{iNy} + e^{-iNy}}{2} + \frac{e^{i(N+1)y} + e^{-i(N+1)y}}{2} + \frac{e^{iy} + e^{-iy}}{2} \sum_{n=-N}^N e^{iny} \\
&= \frac{1}{2} \left(e^{iNy} + e^{-iNy} + e^{i(N+1)y} + e^{-i(N+1)y} + \sum_{n=-N}^N e^{i(n+1)y} + \sum_{n=-N}^N e^{i(n-1)y} \right) \\
&= \frac{1}{2} \left(e^{iNy} + e^{-iNy} + e^{i(N+1)y} + e^{-i(N+1)y} + \sum_{n=-(N-1)}^{N+1} e^{iny} + \sum_{n=-(N+1)}^{N-1} e^{iny} \right) \\
&= \frac{1}{2} \left(\sum_{n=-(N+1)}^{N+1} e^{iny} + \sum_{n=-(N+1)}^{N+1} e^{iny} \right) \\
&= \sum_{n=-(N+1)}^{N+1} e^{iny}
\end{aligned}$$

where we have made use of the induction hypothesis and a tour de force of trigonometric identities.

Now, for any $N \in \mathbb{N}$, due to the periodicity of the functions $x \mapsto e^{inx}$ for $n \neq 0$ and their antiderivatives,

$$\frac{1}{2\pi} \int_{\mathbb{T}} D_N(y) dy = \frac{1}{2\pi} \int_{\mathbb{T}} \sum_{n=-N}^N e^{iny} dy = \frac{1}{2\pi} \int_{\mathbb{T}} e^{i0 \cdot y} dy = \frac{1}{2\pi} \int_{[-\pi, \pi]} 1 dy = 1.$$

Finally, by a quick examination, it is clear the $D_N(y)$ is an even function. Consequently

$$1 = 2 \left(\frac{1}{2\pi} \int_{[0, \pi]} D_N(y) dy \right) = 2 \left(\frac{1}{2\pi} \int_{[-\pi, 0]} D_N(y) dy \right)$$

from which the final result follows. \square

The Dirichlet kernels D_5 , D_{10} and D_{20} are illustrated in Figure 2.




Figure 2: The graphs of D_5, D_{10} and D_{20} .

3.5 Convolutions

Given two functions $f, g \in R(\mathbb{T})$, we define their convolution $f * g : \mathbb{R} \rightarrow \mathbb{C}$ by

$$(f * g)(x) = \frac{1}{2\pi} \int_{\mathbb{T}} f(x - y)g(y) dy = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x - y)g(y) dy$$

for $x \in \mathbb{R}$. The convolution operation is a way to combine information about the functions f and g to produce a new function (which will often be nicer) called $f * g$. Let's consider an example.

4 All things Fourier

4.1 The L^2 theory

In the last subsection, we showed that continuous 2π -periodic functions can be approximated uniformly by trigonometric polynomials. In this subsection, we turn our focus to a larger class of functions, 2π -periodic and Riemann integrable functions, and a completely different mode of approximation, the L^2 approximation. It is my hope that you will see that the L^2 theory, outlined in this subsection, is the cleanest and best-adapted for approximation by trigonometric polynomials. In fact, convergence of Fourier series in L^2 will happen even when other forms of convergence fail.

Let $f : \mathbb{R} \rightarrow \mathbb{C}$. We say that f is Riemann integrable on an interval $[a, b]$ if its real and imaginary parts are Riemann integrable on $[a, b]$. In this case, we define the Riemann integral of f (a complex-valued function) on $[a, b]$ by

$$\int_{[a,b]} f = \int_{[a,b]} f(x) dx = \int_{[a,b]} \operatorname{Re}(f)(x) dx + i \int_{[a,b]} \operatorname{Im}(f)(x) dx.$$

Using the properties of the Riemann integral established for real-valued functions, it is easy to check that the Riemann integral, defined here for complex-valued functions, is also linear.

Now, given a 2π -periodic function $f : \mathbb{R} \rightarrow \mathbb{C}$, we say that f is Riemann integrable on \mathbb{T} if f is Riemann integrable on $[-\pi, \pi]$ and in this case we write

$$\int_{\mathbb{T}} f = \int_{\mathbb{T}} f(x) dx = \int_{[-\pi, \pi]} f(x) dx.$$

When the context is clear, we will simply say that f is Riemann integrable. The set of all such functions will henceforth be denoted by $R(\mathbb{T})$. It is clear that $C(\mathbb{T}) \subseteq R(\mathbb{T})$.

Given $f, g \in R(\mathbb{T})$, we define the $L^2 = L^2(\mathbb{T})$ inner product by

$$\langle f, g \rangle = \frac{1}{2\pi} \int_{\mathbb{T}} f(x) \overline{g(x)} dx.$$

You should verify that $\langle \cdot, \cdot \rangle$ satisfies all of the properties of an inner product except positive definiteness. In particular, $\langle f, f \rangle \geq 0$ for all $f \in R(\mathbb{T})$ and so it makes sense to define the corresponding L^2 norm by

$$\|f\|_{L^2(\mathbb{T})} = \sqrt{\langle f, f \rangle} = \left(\frac{1}{2\pi} \int_{\mathbb{T}} |f(x)|^2 dx \right)^{1/2};$$

this is also called the root-mean-square norm. We will often use the shorthand $\|f\|_2$ for $\|f\|_{L^2(\mathbb{T})}$.

As a quick remark, I should point out that $\|\cdot\|_2$ isn't a bona fide norm on the set $R(\mathbb{T})$. It satisfies all of the properties of a norm except positive definiteness. To see that positive definiteness fails, consider the function

$$f(x) = \begin{cases} 1 & \text{if } x \in 2\pi\mathbb{Z} \\ 0 & \text{otherwise.} \end{cases}$$

It is clear that $f \in R(\mathbb{T})$. Further,

$$\|f\|_2^2 = \frac{1}{2\pi} \int_{[-\pi, \pi]} f(x)^2 dx = 0$$

however f is not the zero function. As it turns out $\|\cdot\|_2$ does become a norm on the Lebesgue space $L^2(\mathbb{T})$ which we do not discuss here. In fact, $L^2(\mathbb{T})$ is a complete (as a metric space) inner product space, such a space is called a Hilbert space. This is standard material in a course on measure theory.

Let us now observe that, for each $n, m \in \mathbb{Z}$,

$$\int_{\mathbb{T}} e^{inx} \overline{e^{imx}} dx = \int_{\mathbb{T}} e^{inx - imx} dx = \int_{\mathbb{T}} e^{i(n-m)x} dx = \int_{[-\pi, \pi]} \cos((n-m)x) dx + i \int_{[-\pi, \pi]} \sin((n-m)x) dx.$$

Consequently,

$$\langle e^{inx}, e^{imx} \rangle = \frac{1}{2\pi} \int_{\mathbb{T}} e^{inx} \overline{e^{imx}} dx = \begin{cases} 1 & \text{if } n = m \\ 0 & \text{if } n \neq m. \end{cases}$$

In view of the preceding calculation, the collection of functions $(e^{inx})_{n \in \mathbb{Z}}$ is called an *orthonormal system*. Let's make a further observation, given $P \in \mathcal{P}(\mathbb{T})$ of the form

$$P(x) = \sum_{n=-N}^N c_n e^{inx},$$

for each $-N \leq m \leq N$,

$$\langle P, e^{imx} \rangle = \sum_{n=-N}^N c_n \langle e^{inx}, e^{imx} \rangle = c_m$$

and $\langle P, e^{imx} \rangle = 0$ whenever $|m| > N$. In this way, we find a way to find the coefficients of a trigonometric polynomial by simply integrating against the elements of the system (e^{inx}) . This is analogous to the way that the coefficients of an analytic function can be computed by taking derivatives. Taking our cues from the above computation, we make a definition.

Definition 4.1. Let $f \in R(\mathbb{T})$. For each $n \in \mathbb{Z}$, define

$$\hat{f}(n) = \langle f, e^{inx} \rangle = \frac{1}{2\pi} \int_{\mathbb{T}} f(x) e^{-inx} dx.$$

The collection of complex numbers $(\hat{f}(n))_{n \in \mathbb{Z}}$ are called the *Fourier coefficients* of f . Considered as a formal series, the series

$$\sum_{n \in \mathbb{Z}} \hat{f}(n) e^{inx}$$

is called the *Fourier series* for f and we write

$$f(x) \sim \sum_{n \in \mathbb{Z}} \hat{f}(n) e^{inx}$$

to indicate that the right hand side is the Fourier series for f .

In view of the discussion preceding the definition, we see that computing the Fourier series corresponding to a trigonometric polynomial P returns the polynomial P itself. Do you see an analogue with Taylor series here? Throughout the rest of this section, we begin to analyze the ways in which the Fourier series for a function f converges. As you will see, if the Fourier series is to converge, it will converge back to f . To this end, we will start talking about partial sums.

Let $f \in R(\mathbb{T})$ and $(\hat{f}(n))_{n \in \mathbb{Z}}$ be the Fourier coefficients of f . For each $N \in \mathbb{N}$, the N th order trigonometric polynomial

$$S_N(x) = \sum_{n=-N}^N \hat{f}(n) e^{inx}$$

defined for $x \in \mathbb{R}$ is called the N th partial sum of the Fourier series $\sum \hat{f}(n) e^{inx}$. Our first main result of the subsection says that, of all N th order trigonometric polynomials, S_N is the best root-mean-square approximation to f .

Theorem 4.2. Let $f \in R(\mathbb{T})$ and let $(\hat{f}(n))_{n \in \mathbb{Z}}$ be its Fourier coefficients. Given $N \in \mathbb{N}$ let $S_N(x)$ be the N th order partial sum of the Fourier series for f and let $P_N \in \mathcal{P}(\mathbb{T})$ be another (possibly different) N th order trigonometric polynomial of the form

$$P_N(x) = \sum_{n=-N}^N c_n e^{inx}.$$

Then

$$\|f - S_N\|_2 \leq \|f - P_N\|_2$$

where equality holds if and only if $c_n = \hat{f}(n)$ for all n .

Proof. We first recall the basic linearity properties of the inner product: For $u, v, w \in R(\mathbb{T})$, and $a \in \mathbb{C}$,

$$\langle u + v, w \rangle = \langle u, w \rangle + \langle v, w \rangle, \quad \langle u, v + w \rangle = \langle u, v \rangle + \langle u, w \rangle$$

and

$$\langle au, v \rangle = a\langle u, v \rangle, \quad \langle u, av \rangle = \bar{a}\langle u, v \rangle.$$

By virtue of these properties, we can write

$$\begin{aligned} \|f - P_N\|_2^2 &= \langle f - P_N, f - P_N \rangle = \langle f, f \rangle - \langle f, P_N \rangle - \langle P_N, f \rangle + \langle P_N, P_N \rangle \\ &= \|f\|_2^2 - \sum_{n=-N}^N \bar{c}_n \langle f, e^{inx} \rangle - \sum_{n=-N}^N c_n \langle e^{inx}, f \rangle + \sum_{n=-N}^N \sum_{m=-N}^N c_n \bar{c}_m \langle e^{inx}, e^{imx} \rangle \\ &= \|f\|_2^2 - \sum_{n=-N}^N \hat{f}(n) \bar{c}_n - \sum_{n=-N}^N c_n \overline{\hat{f}(n)} + \sum_{n=-N}^N c_n \bar{c}_n \\ &= \|f\|_2^2 - \sum_{n=-N}^N |\hat{f}(n)|^2 + \sum_{n=-N}^N |c_n - \hat{f}(n)|^2 \end{aligned}$$

where we have used the fact that $\langle e^{inx}, f \rangle = \overline{\langle f, e^{inx} \rangle} = \overline{\hat{f}(n)}$. Obviously, making the above computation when $c_n = \hat{f}(n)$ yields

$$\|f - S_N\|_2^2 = \|f\|_2^2 - \sum_{n=-N}^N |\hat{f}(n)|^2 \tag{2}$$

and therefore

$$\|f - P_N\|_2^2 = \|f - S_N\|_2^2 + \sum_{n=-N}^N |c_n - \hat{f}(n)|^2$$

from which we observe that

$$\|f - S_N\|_2 \leq \|f - P_N\|_2$$

with equality if and only if $c_n = \hat{f}(n)$ for all n . □

Theorem 4.3 (Bessel's inequality). *Let $f \in R(\mathbb{T})$ and let $(\hat{f}(n))_{n \in \mathbb{Z}}$ be the Fourier coefficients of f . Then*

$$\sum_{n \in \mathbb{Z}} |\hat{f}(n)|^2 \leq \|f\|_2^2 = \frac{1}{2\pi} \int_{\mathbb{T}} |f(x)|^2 dx.$$

In particular, the series

$$\sum_{n \in \mathbb{Z}} |\hat{f}(n)|^2 = \lim_{N \rightarrow \infty} \sum_{n=-N}^N |\hat{f}(n)|^2$$

converges.

Proof. In view of (2), we have

$$\sum_{n=-N}^N |\hat{f}(n)|^2 \leq \|f\|_2^2$$

for all $N \in \mathbb{N}$. The desired result follows by taking the limit of the left hand side as $N \rightarrow \infty$ and noting that the partial sums, whose summands are all non-negative, form a non-decreasing sequence of non-negative numbers. □

Corollary 4.4 (The Riemann-Lebesgue lemma). *Let $f \in R(\mathbb{T})$ and let $(\hat{f}(n))_{n \in \mathbb{Z}}$ be the Fourier coefficients of f . Then*

$$\lim_{n \rightarrow \infty} \hat{f}(n) = \frac{1}{2\pi} \int_{\mathbb{T}} f(x) e^{-inx} dx = 0$$

and

$$\lim_{n \rightarrow \infty} \hat{f}(-n) = \frac{1}{2\pi} \int_{\mathbb{T}} f(x) e^{inx} dx = 0.$$

Proof. The convergence of the series $\sum_{n=-\infty}^{\infty} |\hat{f}(n)|^2$ implies that the summands for sufficiently large and largely negative n converge to zero. \square

Corollary 4.5 (A sharper form of the Riemann-Lebesgue lemma). *Let $[a, b] \subseteq [-\pi, \pi]$ and suppose that f is Riemann integrable on $[a, b]$. Then*

$$\lim_{n \rightarrow \infty} \frac{1}{2\pi} \int_{[a,b]} f(x) \cos(nx) dx = 0$$

and

$$\lim_{n \rightarrow \infty} \frac{1}{2\pi} \int_{[a,b]} f(x) \sin(nx) dx = 0.$$

Proof. Given f as above, consider $g \in R(\mathbb{T})$ defined by

$$g(x) = \begin{cases} f(x) & x \in [a, b] \\ 0 & x \in [-\pi, \pi] \setminus [a, b] \end{cases}$$

and extended periodically to \mathbb{R} . Then, by the Riemann-Lebesgue lemma applied to g , we have

$$\lim_{n \rightarrow \infty} \frac{1}{2\pi} \int_{[a,b]} f(x) e^{-inx} dx = \lim_{n \rightarrow \infty} \frac{1}{2\pi} \int_{[-\pi, \pi]} g(x) e^{-inx} dx = \lim_{n \rightarrow \infty} \hat{g}(n) = 0$$

and

$$\lim_{n \rightarrow \infty} \frac{1}{2\pi} \int_{[a,b]} f(x) e^{inx} dx = \lim_{n \rightarrow \infty} \frac{1}{2\pi} \int_{[-\pi, \pi]} g(x) e^{inx} dx = \lim_{n \rightarrow \infty} \hat{g}(-n) = 0.$$

Consequently,

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{1}{2\pi} \int_{[a,b]} f(x) \cos(nx) dx &= \lim_{n \rightarrow \infty} \frac{1}{2\pi} \int_{[a,b]} f(x) \left(\frac{e^{inx} + e^{-inx}}{2} \right) dx \\ &= \frac{1}{2} \left(\lim_{n \rightarrow \infty} \frac{1}{2\pi} \int_{[a,b]} f(x) e^{inx} dx + \lim_{n \rightarrow \infty} \frac{1}{2\pi} \int_{[a,b]} f(x) e^{-inx} dx \right) \\ &= 0. \end{aligned}$$

The proof that $\lim_{n \rightarrow \infty} (2\pi)^{-1} \int_{[a,b]} f(x) \sin(nx) dx = 0$ is similar. \square

Our next result shows that the Fourier series for f converges to f with respect to the $L^2(\mathbb{T})$ norm. The result also shows that Bessel's inequality is, in fact, an equality. We first need the following simple lemma that you will prove in your Homework 7.

Lemma 4.6. *Let $f \in \mathbb{R}(\mathbb{T})$. For any $\epsilon > 0$, there exists $g \in C(\mathbb{T})$ such that*

$$\|f - g\|_2 < \epsilon.$$

Theorem 4.7 (Parseval’s Theorem). *Let $f \in R(\mathbb{T})$ and let $(\hat{f}(n))_{n \in \mathbb{Z}}$ the Fourier coefficients of f . For each natural number N , denote by S_N the N th partial sum of the Fourier series for f . Then*

$$\lim_{N \rightarrow \infty} \|f - S_N\|_2 = 0;$$

this is the statement that the Fourier series $\sum_{n \in \mathbb{Z}} \hat{f}(n)e^{inx}$ converges to f with respect to the L^2 norm. Further

$$\|f\|_2^2 = \sum_{n \in \mathbb{Z}} |\hat{f}(n)|^2.$$

Proof. Let $\epsilon > 0$ and by an appeal to the lemma choose $h \in C(\mathbb{T})$ for which $\|f - h\|_2 < \epsilon/2$. Now, in view of Theorem ??, Let $P \in \mathcal{P}(\mathbb{T})$ be a trigonometric polynomial of the form

$$P(x) = \sum_{n=-N}^N c_n e^{inx}$$

for which

$$|h(x) - P(x)| < \epsilon/2$$

for all $x \in \mathbb{R}$. From this it follows immediately that

$$\|f - P\|_2 \leq \|f - h\|_2 + \|h - P\|_2 < \epsilon/2 + \left(\frac{1}{2\pi} \int_{[-\pi, \pi]} (\epsilon/2)^2 dx \right)^{1/2} = \epsilon.$$

Now, for any $M \geq N$, the M th partial sum of the Fourier series for f , S_M can be compared with the trigonometric polynomial P (which can trivially be thought of as of M th degree by taking its coefficients to be zero for $M \leq |n| > N$). Thus, in view of Theorem 4.2

$$\|f - S_M\|_2 \leq \|f - P\|_2 < \epsilon.$$

Hence, for every $\epsilon > 0$, there exists and $N \in \mathbb{N}$ such that, for every $M \geq N$, $\|f - S_M\|_2 < \epsilon$ and therefore

$$\lim_{N \rightarrow \infty} \|f - S_N\|_2 = 0.$$

With this, (2) gives

$$\lim_{N \rightarrow \infty} \sum_{n=-N}^N |\hat{f}(n)|^2 = \|f\|_2^2 - \lim_{N \rightarrow \infty} \|f - S_N\|_2^2 = \|f\|_2^2 - 0$$

from which the desired result follows immediately. □

Example 1

Consider the so-called sawtooth function defined by

$$f(x) = x \quad -\pi < x \leq \pi$$

and extended 2π -periodically to \mathbb{R} . The graph of f is illustrated in Figure 3 (the vertical lines are not part of the graph; they are inserted automatically by Matlab).




Figure 3: The graph of f for $-3\pi < x \leq 3\pi$.

Let's compute the Fourier coefficients of f . For $n = 0$, we have

$$\hat{f}(0) = \frac{1}{2\pi} \int_{\mathbb{T}} f(x) e^{-i \cdot 0 \cdot x} dx = \frac{1}{2\pi} \int_{\mathbb{T}} x dx = \frac{1}{2\pi} \int_{[-\pi, \pi]} x dx = 0.$$

For $n \neq 0$,

$$\begin{aligned} 2\pi \hat{f}(n) &= \int_{\mathbb{T}} x e^{-inx} dx \\ &= \int_{[-\pi, \pi]} x \cos(-nx) dx + i \int_{[-\pi, \pi]} x \sin(-nx) dx \\ &= \int_{[-\pi, \pi]} x \cos nx dx - i \int_{[-\pi, \pi]} x \sin nx dx \\ &= -i \left(\frac{-1}{n} x \cos nx \Big|_{-\pi}^{\pi} - \frac{-1}{n} \int_{[-\pi, \pi]} \cos nx dx \right) \\ &= \frac{i}{n} (\pi \cos n\pi - (-\pi \cos(-n\pi))) - 0 \\ &= \frac{2\pi i}{n} \cos n\pi = \frac{2\pi i}{n} (-1)^n \end{aligned}$$

where we have integrated by parts and used (heavily) the periodicity and odd and even properties of sine/cosine. Consequently,

$$\hat{f}(n) = \begin{cases} \frac{i(-1)^n}{n} & n \neq 0 \\ 0 & n = 0. \end{cases}$$

Observe that

$$\sum_{n \in \mathbb{Z}} |\hat{f}(n)|^2 = \sum_{n \in \mathbb{Z} \setminus \{0\}} \frac{1}{n^2} = 2 \sum_{n=1}^{\infty} \frac{1}{n^2}.$$

Also,

$$\|f\|_2^2 = \frac{1}{2\pi} \int_{\mathbb{T}} |f(x)|^2 dx = \frac{1}{2\pi} \int_{[-\pi, \pi]} x^2 dx = \frac{1}{2\pi} \frac{x^3}{3} \Big|_{-\pi}^{\pi} = \frac{2\pi^3}{6\pi} = \frac{\pi^2}{3}$$

and so, by an application of Parseval’s theorem, we obtain

$$\frac{\pi^2}{3} = \|f\|_2^2 = \sum_{x \in \mathbb{Z}} |\hat{f}(n)|^2 = 2 \sum_{n=1}^{\infty} \frac{1}{n^2}.$$

It was long understood that the series $\sum_{n=1}^{\infty} 1/n^2$ converged. The Basel problem, posed by Pietro Mengoli around 1644, asks: What is the exact value of this series? In 1734, a mathematician named Leonhard Euler showed that the value of the series is exactly $\pi^2/6$. The solution made Euler instantly famous. By simply dividing the previous equation by 2 we obtain Euler’s result (Theorem 4.8 below). We note that our approach (via Parseval’s Theorem) is completely different than that of Euler. This should not be surprising as Fourier series wasn’t discovered for nearly one hundred years after Euler presented his result.

Theorem 4.8 (Euler 1734).

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}.$$

As our attention will soon turn to the investigation of pointwise convergence of Fourier series, let’s continue our Fourier series analysis of the Sawtooth function. As we will see, this example contains a surprising number of the strange phenomena/pathologies found commonplace in the study of Fourier series, including the Gibb’s phenomenon. For each N , the N th Fourier polynomial for f is given by

$$\begin{aligned} S_N(x) &= \sum_{n=-N}^N \hat{f}(n)e^{inx} = \hat{f}(0) + \sum_{n=1}^N \hat{f}(n)e^{inx} + \sum_{n=1}^N \hat{f}(-n)e^{-inx} \\ &= \sum_{n=1}^N \frac{i(-1)^n}{n} e^{inx} + \frac{i(-1)^{-n}}{-n} e^{-inx} = \sum_{n=1}^N \frac{(-1)^n}{n} i(e^{inx} - e^{-inx}) \\ &= \sum_{n=1}^N \frac{(-1)^n}{n} i(2i \sin nx) = \sum_{n=1}^N \frac{2(-1)^{n+1}}{n} \sin nx \end{aligned}$$

for $x \in \mathbb{R}$. The Fourier polynomials of degrees 1, 2, 3, 4, 5 and 40 are illustrated in Figure 4 for $-3\pi < x \leq 3\pi$. The reader should note that the partial sums appear to be converging nicely except for the overshoot near the points of discontinuity at $x = \pm 3\pi, \pm \pi$; this is the Gibb’s phenomenon.




Figure 4: The graphs of f and P_n for $n = 1, 2, 3, 4, 5, 40$.

4.2 The pointwise and uniform theory theory

As we saw in the last subsection, the Fourier series for a function $f \in R(\mathbb{T})$ converges to f with respect to the L^2 norm. In this subsection, we investigate the same question from the perspective of pointwise (and uniform)

convergence; as we have seen, both notions are stronger than L^2 convergence. When originally investigating Fourier series in its connection to the theory of heat, Jean-Baptiste Fourier claimed that, given $f \in C(\mathbb{T})$, the Fourier series $\sum_{n \in \mathbb{N}} \hat{f}(n)e^{inx}$ converges to $f(x)$ for all $x \in \mathbb{R}$. As it turns out, this isn't true.

Theorem 4.9 (Du Bois-Reymond, 1873). *There exists $f \in C(\mathbb{T})$ whose Fourier series $\sum_{n \in \mathbb{Z}} \hat{f}(n)e^{inx}$ diverges at a point $x \in (-\pi, \pi]$. This means specifically that, for some $x \in (-\pi, \pi]$, the limit*

$$\lim_{N \rightarrow \infty} \sum_{n=-N}^N \hat{f}(n)e^{inx}$$

does not exist.

Following the discovery of this result, a frantic search began in mathematics to find precise conditions on a function f which would guarantee that its Fourier series converged. Throughout this search, it was discovered that the Riemann integral was insufficient for the needed purposes of investigation and this helped lead to a complete revolution in mathematics. Out of the revolution emerged the Lebesgue integral and Lebesgue's theory of integration. Any serious investigation of Fourier series (which ours is unfortunately not) requires one to understand the Lebesgue integral. In 1926 a brilliant young mathematician named Andre Kolmogorov showed that things were much worse than had been previously thought (and had been shown by Du Bois-Reymond).

Theorem 4.10 (Kolmogorov 1926). *There exists a Lebesgue integrable function f (we say $f \in L^1(\mathbb{T})$) whose Fourier series diverges at every point.*

We now begin our investigation of the uniform convergence of Fourier series.

Lemma 4.11. *Let $f, g \in C(\mathbb{T})$, if $\widehat{f-g}(n) = \hat{f}(n) - \hat{g}(n)$ for all n , then $f = g$, i.e., $f(x) = g(x)$ for all $x \in \mathbb{R}$.*

Proof. As you show in your homework, $\widehat{f-g}(n) = \hat{f}(n) - \hat{g}(n)$ for all $n \in \mathbb{Z}$, i.e., the map $f \mapsto \hat{f}$ is linear. Thus, $\widehat{f-g}(n) = 0$ for all $n \in \mathbb{Z}$ and so, by an appeal to Parseval's theorem,

$$\|f - g\|_2^2 = \sum_{n \in \mathbb{Z}} |\widehat{f-g}(n)|^2 = 0.$$

Consequently,

$$\int_{\mathbb{T}} |f(x) - g(x)|^2 dx = 2\pi \|f - g\|_2^2 = 0.$$

It now follows, by the result I prove in class during Week 6, that $f(x) - g(x) = 0$ for all $x \in [-\pi, \pi]$. Because f and g are 2π periodic, we conclude that $f(x) = g(x)$ for all $x \in \mathbb{R}$. \square

With the properties of the Dirichlet kernel established in the preceding proposition, we have our first (truly) pointwise result.

Theorem 4.12. *Let $f \in R(\mathbb{T})$ and assume that the $f = u + iv$ is piecewise differentiable, i.e., its real and imaginary parts u and v are continuously differentiable on $[-\pi, \pi]$ except possibly at a finite number of points where u, v, u' and v' have (at worse) removable or jump discontinuities. For any $x_0 \in \mathbb{R}$, define*

$$f(x_0^+) = \lim_{x \rightarrow x_0; x > x_0} f(x) \quad \text{and} \quad f(x_0^-) = \lim_{x \rightarrow x_0; x < x_0} f(x).$$

Then, at each $x_0 \in \mathbb{R}$, the Fourier series for f converges and

$$\frac{f(x_0^+) + f(x_0^-)}{2} = \sum_{n \in \mathbb{Z}} \hat{f}(n)e^{inx_0} = \lim_{N \rightarrow \infty} \sum_{n=-N}^N \hat{f}(n)e^{inx_0}.$$

In particular, if f is continuous at x_0 (or has a removable discontinuity at x_0),

$$f(x_0) = \sum_{n \in \mathbb{Z}} \hat{f}(n)e^{inx_0};$$

otherwise, the Fourier series for f converges to the average of the left and right limits of f at x_0 .

Before proving the theorem, let's illustrate its conclusion by revisiting the sawtooth function and its Fourier series.

Example 2

We recall the sawtooth function defined by $f(x) = x$ for $-\pi < x \leq \pi$ and extended periodically to \mathbb{R} . We previously studied this function in Example 4.1 and computed its Fourier series. We found

$$f(x) \sim \sum_{n \in \mathbb{Z} \setminus \{0\}} \frac{i(-1)^n}{n} e^{inx} = \sum_{n=1}^{\infty} \frac{2(-1)^{n+1}}{n} \sin nx$$

Now, by a straightforward computation, f is differentiable on the open set $\mathbb{R} \setminus \{(2k+1)\pi : k \in \mathbb{Z}\}$ (consisting of the entire real line except for the *breakpoints* $\pm\pi, \pm3\pi, \pm5\pi, \dots$) and, on this set, $f'(x) = 1$. Consequently, f is piecewise differentiable and so we may apply the theorem.

For $-\pi < x < \pi$, f is continuous and by virtue of the theorem we conclude that the Fourier series for f converges to $f(x) = x$ on the open set $(-\pi, \pi)$. At $x = \pi$, f has a discontinuity. Here we have

$$f(\pi_-) = \lim_{x \rightarrow \pi^-; x < \pi} f(x) = \lim_{x \rightarrow \pi^-; x < \pi} x = \pi$$

and

$$f(\pi_+) = \lim_{x \rightarrow \pi^+; x > \pi} f(x) = \lim_{x \rightarrow \pi^+; x > \pi} (x - 2\pi) = \pi - 2\pi = -\pi.$$

Consequently,

$$\sum_{n \in \mathbb{Z}} \hat{f}(n) e^{in\pi} = \lim_{N \rightarrow \infty} S_N(\pi) = \frac{1}{2}(\pi + -\pi) = 0;$$

in fact, this result holds at all the breakpoints $\pm\pi, \pm3\pi, \dots$. Appealing to the full scope of the theorem (or simply noting that the above conclusion extends by periodicity), we have

$$\sum_{n \in \mathbb{Z}} \hat{f}(n) e^{inx} = \lim_{N \rightarrow \infty} S_N(x) = \begin{cases} (x - 2\pi k) & (2k-1)\pi < x < (2k+1)\pi, k \in \mathbb{Z} \\ 0 & x = (2k+1)\pi, k \in \mathbb{Z} \end{cases}.$$

In other words, the Fourier series for f converges to f pointwise on the open set $\mathbb{R} \setminus \{(2k+1)\pi : k \in \mathbb{Z}\}$ and it converges to 0 elsewhere. We note that the series does not converge uniformly for, if this were the case, f would be continuous. This convergence is illustrated in the following Figure 5.




Figure 5: f and S_{40}

We now prove the theorem.

Proof of Theorem 4.12. Our aim is to show that

$$\lim_{N \rightarrow \infty} \left(S_N(x_0) - \frac{f(x_0^+) + f(x_0^-)}{2} \right) = 0.$$

In view of Proposition 3.24, we have

$$\begin{aligned} \left(S_n(x_0) - \frac{f(x_0^+) + f(x_0^-)}{2} \right) &= \frac{1}{2\pi} \int_{\mathbb{T}} f(x_0 - y) D_N(y) dy - \left(f(x_0^+) \frac{1}{2} + f(x_0^-) \frac{1}{2} \right) \\ &= \frac{1}{2\pi} \int_{[-\pi, 0]} f(x_0 - y) D_N(y) dy + \frac{1}{2\pi} \int_{[0, \pi]} f(x_0 - y) D_N(y) dy \\ &\quad - \left(f(x_0^+) \frac{1}{2\pi} \int_{[-\pi, 0]} D_N(y) dy + f(x_0^-) \frac{1}{2\pi} \int_{[0, \pi]} D_N(y) dy \right) \\ &= \frac{1}{2\pi} \int_{[-\pi, 0]} (f(x_0 - y) - f(x_0^+)) D_N(y) dy + \frac{1}{2\pi} \int_{[0, \pi]} (f(x_0 - y) - f(x_0^-)) D_N(y) dy \\ &=: I_+(N) + I_-(N). \end{aligned}$$

We consider the integrals I_+ and I_- . By virtue of Proposition 3.24,

$$\begin{aligned}
I_+(N) &= \frac{1}{2\pi} \int_{[-\pi, 0]} (f(x_0 - y) - f(x_0^+)) \frac{\sin((N + 1/2)y)}{\sin(y/2)} dy \\
&= \frac{1}{2\pi} \int_{[0, \pi]} (f(x_0 + y) - f(x_0^+)) \frac{\sin((N + 1/2)y)}{\sin(y/2)} dy \\
&= \frac{1}{2\pi} \int_{[0, \pi]} (f(x_0 + y) - f(x_0^+)) \left(\cos(Ny) + \frac{\cos(y/2) \sin(Ny)}{\sin(y/2)} \right) dy \\
&= \frac{1}{2\pi} \int_{[0, \pi]} (f(x_0 + y) - f(x_0^+)) \cos(Ny) dy + \frac{1}{2\pi} \int_{[0, \pi]} \left(\frac{f(x_0 + y) - f(x_0^+)}{y} \right) \left(2(y/2) \frac{\cos(y/2)}{\sin(y/2)} \right) \sin(Ny) dy \\
&= \frac{1}{2\pi} \int_{[0, \pi]} g(y) \cos(Ny) dy + \frac{1}{2\pi} \int_{[0, \pi]} h(y) \sin(Ny) dy
\end{aligned}$$

where

$$g(y) = f(x_0 + y) - f(x_0^+)$$

and

$$h(y) = \left(\frac{f(x_0 + y) - f(x_0^+)}{y} \right) \left(2(y/2) \frac{\cos(y/2)}{\sin(y/2)} \right).$$

It is clear that $g \in R(\mathbb{T})$ and thus, by Corollary 4.5,

$$\lim_{N \rightarrow \infty} \frac{1}{2\pi} \int_{[0, \pi]} g(y) \cos(Ny) dy = 0.$$

Making the same conclusion concerning h isn't so straightforward. First, given our hypotheses concerning f , it is clear that h is piecewise continuous (continuous except at a finite number of points) on the interval $(0, \pi]$ and so it is Riemann-integrable on every compact subinterval of $(0, \pi]$. We must examine h near $y = 0$ for the only possible impediment for Riemann integrability on $[0, \pi]$ is the behavior of h as $y \rightarrow 0$. First,

$$\lim_{y \rightarrow 0; y > 0} 2(y/2) \frac{\cos(y/2)}{\sin(y/2)} = \lim_{y \rightarrow 0; y > 0} 2 \cos(y/2) \frac{(y/2)}{\sin(y/2)} = 2.$$

Secondly, because f is piecewise differentiable on $[-\pi, \pi]$,

$$\lim_{y \rightarrow 0; y > 0} \frac{f(x_0 + y) - f(x_0^+)}{y} = \lim_{y \rightarrow 0; y > 0} f'(x_0 + y) = f'(x_0^+).$$

We remark that this is really a statement about exchanging limits and derivatives and its validity is far from obvious. A rigorous proof of this limit (which you should attempt), can be seen as an application of Theorem ???. Putting these two result together shows that

$$\lim_{h \rightarrow 0; h > 0} h(y) = 2f'(x_0^+).$$

In particular, h is bounded (and well-behaved) at 0 and so it follows that h is Riemann integrable on $[0, \pi]$. By an application of Corollary 4.5, we conclude that

$$\lim_{N \rightarrow \infty} \frac{1}{2\pi} \int_{[0, \pi]} h(y) \sin(Ny) dy = 0.$$

Consequently,

$$\lim_{N \rightarrow \infty} I_+(N) = 0.$$

By making completely analogous reasoning, it follows that

$$\lim_{N \rightarrow \infty} I_-(N) = 0$$

and therefore

$$\lim_{N \rightarrow \infty} \left(S_N(x_0) - \frac{f(x_0^+) + f(x_0^-)}{2} \right) = \lim_{N \rightarrow \infty} (I_+(N) + I_-(N)) = 0.$$

□

Exercise 11

We've seen three results so far about the convergence of Fourier series, Theorem 3.22 (and related Corollary 3.23), Theorem 4.7 and Theorem 4.12. Under certain hypotheses, each of these theorems gives an affirmative (in a certain sense) to the question: Can a function be expanded in terms of its Fourier series? In this exercise, you are asked to analyze a handful of specific examples through the lens of these three theorems. Specifically, for each of the functions f below, do the following:

- Determine if the given function f is a member of $R(\mathbb{T})$, $C^m(\mathbb{T})$ for any $m \geq 0$ and/or the set of piecewise differentiable functions on \mathbb{T} .
- Compute the Fourier coefficients of f .
- Compute and simplify the Fourier series for f .
- In view of all of the results discussed above, make the strongest statement you can about the convergence of the Fourier series of f . For instance, does it converge pointwise? If so, to what? Does it converge uniformly? If so, to what? Does it converge with respect to the L^2 norm? Please give precise statements.
- If you concluded that $f \in R(\mathbb{T})$ (the hypothesis of Theorem 4.7), you should expect that

$$\|f\|_2^2 = \sum_{n \in \mathbb{Z}} |\hat{f}(n)|^2.$$

Compute both sides of this identity (Parseval's identity). As our example of the sawtooth function gave us the solution to Basel's problem, $\sum_{n=1}^{\infty} 1/n^2 = \pi^2/6$, what does Parseval's identity give you here?

The following is the list of functions to be analyzed. If the rule of the function f is only given on the interval $(-\pi, \pi]$, the function should be assumed to be 2π -periodic on \mathbb{R} (and so the rule on $(-\pi, \pi]$ determines the function completely).

1.

$$f(x) = |x| \quad \text{for } x \in (-\pi, \pi].$$

2. For $\alpha \in \mathbb{R}$,

$$f(x) = \cos(\alpha x) \quad \text{for } x \in (-\pi, \pi].$$

You should expect a discrepancy between the cases $\alpha \in \mathbb{Z}$ and $\alpha \in \mathbb{R} \setminus \mathbb{Z}$.

3.

$$f(x) = \begin{cases} 1 & x \in (0, \pi] \\ 0 & x \in (-\pi, 0] \end{cases}.$$

4.

$$f(x) = \begin{cases} x(\pi - x) & x \in (0, \pi] \\ x(\pi + x) & x \in (-\pi, 0] \end{cases}.$$

Theorem 4.12 is the strongest result about the convergence of Fourier series we will prove in this course. I hope that you find the result satisfying, it encompasses most of the functions that you know about and can write down. The

Nobel prize-winning physicist, Richard Feynman, was quite happy with this results (and ones like it) when made the following statement: “The mathematicians have shown, for a wide class of functions, in fact for all that are of interest to physicists, that if we can do the integrals we will get back $f(t)$.” He made this statement in the 1960’s in his famous lecture series at Caltech, just before Lennart Carleson completely solved the problem and determined the exact class of functions representable by their Fourier series [?, ?]. Carleson’s result, now known as Carleson’s theorem, was a long standing conjecture known as Luzin’s conjecture. For your cultural benefit, I will state it here; I’ll first need to make a definition.

Definition 4.13. For any interval $I = [a, b] \subseteq \mathbb{R}$, we define

$$\ell(I) = b - a$$

to be the length of I . Now, for any subset E of \mathbb{R} , we say that E is a set of measure zero (or a null set) if, for every $\epsilon > 0$, there is an infinite collection of intervals $\{I_n\}$ such that

$$E \subseteq \bigcup_{n=1}^{\infty} I_n$$

and

$$\sum_{n=1}^{\infty} \ell(I_n) < \epsilon.$$

You should think of a set of measure zero as an extremely small set. For instance, any countable (and hence finite) collection of points is of measure zero. There are, however, uncountable sets of measure zero, an important example of which is the Cantor set.

Exercise 12

In this exercise, you verify the claim made in the penultimate sentence. Prove the following statement:

If $E \subseteq \mathbb{R}$ is countable, i.e., there exists a surjection (onto function) $\phi : \mathbb{N} \rightarrow E$, then E has measure zero.

Hint: Surround each point $x_k \in E$ by an interval whose length is $\epsilon/2^{k+1}$.

Using this notion of small sets we can state Carleson’s theorem in the context of Riemann integrable functions; the general result is formulated using the Lebesgue integral [?]. Here it is:

Theorem 4.14 (Carleson 1966). For any $f \in R(\mathbb{T})$, there exists a set E of measure zero (which is possibly empty) such that

$$f(x) = \sum_{n \in \mathbb{Z}} \hat{f}(n) e^{inx} = \lim_{N \rightarrow \infty} S_N(x)$$

for all $x \notin E$.

4.3 The Gibb’s phenomenon

In this final subsection, we will study the Gibb’s phenomenon. The Gibb’s phenomenon, named after J. Willard Gibb’s (yes, the free energy Gibbs), describes the pointwise convergence of Fourier series of a function with a jump discontinuity. Instead of working in the general setting, we will study the Gibb’s phenomenon as it occurs when we consider the Fourier series of the sawtooth function. Focusing on this specific case will allow us to very precisely see what’s going on. If you are worried about the general case, I’ll refer you to a very nice discussion by T. W. Körner in which he describes how to extend the results pertaining to this example to the general class of piecewise

differentiable functions in $R(\mathbb{T})$ [?].

So let's return to our favorite example, the sawtooth function f , defined by

$$f(x) = x$$

for all $-\pi < x \leq \pi$ and extended periodically to \mathbb{R} . We recall that

$$f(x) \sim \sum_{n \in \mathbb{Z} \setminus \{0\}} \frac{(-1)^n}{n} e^{inx} = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} \sin(nx) \tag{3}$$

Let's again consider the graph of the Fourier polynomial S_{40} ; this is illustrated in Figure 6.




Figure 6: f and S_{40}

As we discussed in the last subsection, Theorem 4.12 guarantees that

$$\lim_{N \rightarrow \infty} S_N(x) = f(x)$$

for all $x \neq m\pi$ where $m \in \mathbb{Z}$ is odd. We also showed that, at any $x = m\pi$ where $m \in \mathbb{Z}$ is odd, $S_N(x) \rightarrow 0$ as $N \rightarrow \infty$. These two things should appear to be somewhat clear by looking at Figure 6. There is however one thing that should bother you: near (but not at) the breakpoints, $\{\dots, -3\pi, -\pi, \pi, 3\pi, \dots\}$, the graph of S_{40} seems to “overshoot” (or “undershoot”) the graph of f . These are the spikes you see close to the discontinuity in f . This behavior is called the *Gibb’s phenomenon*. Let’s study this behavior more closely at, say, $x = \pi$; Figure 7 shows the graphs of $S_N(x)$ for $N = 25, 26, \dots, 50$.




Figure 7: The graphs of $S_n(x)$ for $n = 25, 26, \dots, 50$ and $f(x)$ for $9\pi/10 \leq x \leq \pi$.

Upon studying Figure 7 closely, we see that the overshoot is moving right as N increases. You might say: Theorem 4.12 guarantees that $S_N(x) \rightarrow f(x) = x$ for all $-\pi < x < \pi$ but, upon looking at the figure, $S_N(x)$ isn't converging to $f(x)$ for x very close to π . So where did we go wrong? The answer is that we haven't gone wrong at all, the apparent discrepancy can be understood by recognizing that pointwise convergence is weaker than convergence in the graph—this is the difference between pointwise convergence and uniform convergence. Remember, that for pointwise convergence, we first select x and ϵ and find a natural number $M = M(\epsilon, x)$ for which

$$|S_N(x) - f(x)| < \epsilon$$

for all $N \geq M$. In the case at hand, we can understand this notion as follows: If I select an $x < \pi$, but as close to π as I want, since the overshoot in the Fourier polynomials are moving to the right, I simply have to wait until they have moved so far right that they've passed x —this will determine N . After this, the Fourier polynomials evaluated at x will get much much closer to $f(x)$. Okay, so now you understand how we still get pointwise convergence. Let's now try to understand the overshoot.

Using the same numbers I've used to make the graphs in Figure 7, I can quantify this overshoot. For each $N \in \mathbb{N}$, denote by

$$M_N = \max_{9\pi/10 \leq x \leq \pi} S_N(x),$$

the maximum of the function $S_N(x)$ near π . We also denote by x_N the unique x near π for which

$$f(x_N) = M_N.$$

The following table shows M_N , M_N/π , x_N and $\pi - \pi/N$ to four decimal places for $n = 25, 30, \dots, 50$.

N	M_N	M_N/π	x_N	$\pi - \pi/N$
25	3.5822	1.1403	3.0204	3.0159
30	3.6020	1.1465	3.0404	3.0369
35	3.6172	1.1514	3.0544	3.0518
40	3.6321	1.1561	3.0654	3.0631
45	3.6433	1.1597	3.0734	3.0718
50	3.6516	1.1624	3.0804	3.0788

Upon looking at the table, we see that, as N increases the x_N 's are close to $\pi - \pi/N$ and the ratio M_N/π grows toward 1.17... So, by following the x at which $S_N(x)$ is maximized, the ratio M_N/π approaches some number A as $N \rightarrow \infty$ (note that π is half the gap of the discontinuity of f at π); this describes the overshoot. The following theorem formalizes it:

Theorem 4.15. *Let f, S_N be as above. Then*

$$\lim_{N \rightarrow \infty} S_N(\pi - \pi/N) = \pi A$$

where

$$A = 1.178979744447216727 \dots$$

Thus, the $S_N(\pi - \pi/N)/\pi$ converges to A (called the Gibb's constant) times half of the gap of the jump discontinuity.

Proof. Using our trigonometric identities, we find that

$$\begin{aligned} S_N(\pi - \pi/N) &= \sum_{n=1}^N \frac{2(-1)^{n+1}}{n} \sin(n\pi - n\pi/N) \\ &= \sum_{n=1}^N \frac{2(-1)^{n+1}}{n} (\sin(n\pi) \cos(n\pi/N) - \sin(n\pi/N) \cos(n\pi)) \\ &= \sum_{n=1}^N \frac{2(-1)^{n+1}}{n} (0 - \sin(n\pi/N) \cos(n\pi)) \\ &= \sum_{n=1}^N \frac{2(-1)^{n+1}}{n} ((-1)^{n+1} \sin(n\pi/N)) \\ &= 2 \sum_{n=1}^N \frac{\sin n\pi/N}{n\pi/N} \frac{\pi}{N}. \end{aligned}$$

You should recognize that

$$\sum_{n=1}^N \frac{\sin n\pi/N}{n\pi/N} \frac{\pi}{N}$$

is a (right) Riemann sum for the integral

$$\int_0^\pi \frac{\sin(x)}{x} dx$$

and because $\sin x/x$ is Riemann integrable on $[0, \pi]$, we immediately conclude that

$$\lim_{N \rightarrow \infty} S_N(\pi - \pi/N) = \lim_{N \rightarrow \infty} 2 \sum_{n=1}^N \frac{\sin n\pi/N}{n\pi/N} \frac{\pi}{N} = 2 \int_0^\pi \frac{\sin x}{x} dx = \pi A$$

where

$$A = \frac{2}{\pi} \int_0^\pi \frac{\sin x}{x} dx.$$

It remains to compute A . Using the power series representation for $\sin x$ about 0, we have

$$\frac{\sin x}{x} = \frac{1}{x} \left(x - \frac{x^3}{3!} + \frac{x^5}{5!} \dots \right) = 1 - \frac{x^2}{3!} + \frac{x^4}{5!} + \dots$$

for $x \in \mathbb{R}$ where we take the left hand side to be 1 when $x = 0$. Using the results of Exercise C, it is easily verified that the above series converges absolutely and uniformly on $[0, \pi]$. By an application of Corollary C.9, we conclude

that

$$\begin{aligned} A &= \frac{2}{\pi} \int_0^\pi \left(1 - \frac{x^2}{3!} + \frac{x^4}{5!} + \dots\right) dx = \frac{2}{\pi} \left(\int_0^\pi 1 dx - \frac{1}{3!} \int_0^\pi x^2 dx + \frac{1}{5!} \int_0^\pi x^4 dx \right) \\ &= \frac{2}{\pi} \left(\pi - \frac{\pi^3}{3 \cdot 3!} + \frac{\pi^5}{5 \cdot 5!} + \dots \right) = 2 - \frac{2\pi^2}{3 \cdot 3!} + \frac{2\pi^4}{5 \cdot 5!} \dots \\ &= 1.178979744447216727\dots \end{aligned}$$

as desired. □

Exercise 13

There is (at least) one function f in Exercise 16 that has a jump discontinuity in the interval $[-\pi, \pi]$. For this example, plot (in any computing program you want) some Fourier polynomials of f . Print out the results and comment on the appearance (or lack thereof) of the Gibb's phenomenon. If you'd like a source of the Matlab file I used to analyze the sawtooth function, let me know and I'll email it to you.

5 The Fourier Transform

A Sups, Infs and all that

In this short section we discuss the essential properties of real numbers that will we need in our study of (Fourier) analysis. Though most of our studies will involve complex-valued things (numbers, functions, etc.), it is essential that we understand some basic objects of mathematical analysis on the real line – these objects provide the foundation upon which the notions of distance and convergence is built.

Definition A.1 (Bounded Sets). *Let A be a subset of real numbers.*

1. We say that the set A is bounded above if there exists a real number M for which

$$x \leq M \quad \text{for all} \quad x \in A.$$

In this case, M is said to be an upper bound for the set A . We also say that A is bounded above by M .

2. We say that the set A is bounded below if there exists a real number N for which

$$N \leq x \quad \text{for all} \quad x \in A.$$

In this case N is said to be a lower bound for the set A . We also say that A is bounded below by N .

3. We say that the set A is bounded if it is bounded above and below.

In light of the following definition, we observe that a bounded set A can be contained inside a “closed” interval, i.e., if A is bounded then there are constants N and M for which $A \subseteq [N, M]$. This is an important topological notion that extends well beyond the real line. The following exercise expands on this.

Exercise 14

Given a real number a and a positive real number r , we define the (open) ball centered at a with radius r as the set

$$B_r(a) = \{x \in \mathbb{R} : |x - a| < r\} = (a - r, a + r).$$

Let A be a set of real numbers. Prove that the following are equivalent.

1. A is bounded.

2. There exists $r > 0$ for which $A \subseteq B_r(0)$.
3. There exists $a \in \mathbb{R}$ and $r > 0$ for which $A \subseteq B_r(a)$.

Hint: Prove the implication (1) \rightarrow (2) \rightarrow (3) \rightarrow (1).

Perhaps the most important notion for (bounded) sets of real numbers is captured by the following definition.

Definition A.2 (Supremum). *Let A be a set of real numbers which is bounded above. Suppose that there exists a number S with the following two properties:*

- i. S is an upper bound of A .*
- ii. For any upper bound M of A , $s \leq M$.*

Then S is called the supremum (or least upper bound) of the set A and we write $S = \sup A$.

The careful reader should note that I've used the definite article ("the") in the above definition. This language seems to imply that, given a set A which is bounded above, that there can be at most one number called the supremum (of A). Let's justify this.

Lemma A.3. *Let A be a set of real numbers which is bounded above and suppose that S_1 and S_2 are numbers which both satisfy Conditions (i) and (ii) of the preceding definition. Then $S_1 = S_2$.*

Proof. In view of our hypothesis, S_1 and S_2 are both upper bounds for A . Thus, given that condition (ii) is satisfied for S_1 , we have $S_1 \leq S_2$ (because S_2 is an upper bound for A). Similarly, given that (ii) is satisfied for S_2 , $S_2 \leq S_1$. Thus, $S_1 \leq S_2$ and $S_2 \leq S_1$ whence $S_1 = S_2$. \square

In light of the above result, to each $A \subseteq \mathbb{R}$ which is bounded above, there is at most one supremum for A . The natural thing to ask is the following. Must such a set A have a supremum at all? Though we will not worry about it in this course, the answer to this question is deeply rooted in the construction of the real numbers. We will therefore take the following for granted.

Theorem A.4 (The completeness axiom, Theorem 1.11 of [?]). *Suppose A is a non-empty subset of \mathbb{R} which is bounded above. Then*

$$S = \sup A$$

exists.

In the case that a non-empty set A is not bounded above, we assume the common convention and write

$$\sup A = \infty.$$

Here (and being consistent with Theorem A.4), $\sup A$ is only understood to exist in an extended sense (as an extended real number). Here we will always assume a clear distinction between real numbers and ∞ . It is still however instructive to have $\sup A$ be able to take the "value" of ∞ when A is unbounded.

It will be useful for us to have a characterization of the supremum of a bounded set A . This condition, which we will make use of often, is the subject of the following proposition.

Proposition A.5. *Let A be a non-empty subset of real numbers which is bounded above and let S be an upper bound of A . Then S is the supremum of A if and only if the following condition is satisfied.*

$$\text{For each } \epsilon > 0, \text{ there is an element } x \in A \text{ for which } x > S - \epsilon.$$

This condition is illustrated in Figure 8.




Figure 8: An illustration of the characterizing condition in Proposition A.5

Proof. Suppose that $S = \sup A$ and let $\epsilon > 0$. We observe that $S - \epsilon < S$. Since S is the supremum (the least upper bound) of A , any number strictly less than S cannot be an upper bound of A and so $S - \epsilon$ cannot be an upper bound of A . Since A is non-empty, we must therefore have some $x \in A$ for which $S - \epsilon < x$. This proves our desired property.

Conversely, assume that S is an upper bound of A satisfying the property that, for all $\epsilon > 0$ there is an element $x \in A$ for which $x > S - \epsilon$. Let M be another upper bound of A . In the case that $M < S$, we set $\epsilon = S - M > 0$ and observe that, by the supposition, there is an $x \in A$ for which $S - \epsilon < x$. We note however that $S - \epsilon = S - (S - M) = M$ and thus there is an $x \in A$ for which $x > M$ showing that M cannot be an upper bound for A , a contradiction. Hence $S \leq M$ and so $S = \sup A$. \square

To see the utility of the proposition above, let us apply the proposition in the case where A is a subinterval of the real line. As you expect, the least upper bound of the interval should be its larger end point. Though these seems obvious, the claim requires some checking which we do in the following example.

Example 1

Let $a < b$ be real numbers and let I be an interval of the form $[a, b]$, $(a, b]$, $[a, b)$ or (a, b) . Then

$$\sup I = b.$$

To prove the statement above, let $\epsilon > 0$ and observe that $b - \epsilon < b$. Let

$$x = \max \left\{ b - \frac{b-a}{2}, b - \frac{\epsilon}{2} \right\}$$

and observe that, by definition,

$$a = b - (b-a) \leq b - \frac{b-a}{2} \leq x < b$$

and

$$b - \epsilon \leq b - \frac{\epsilon}{2} \leq x < b.$$

Thus x is a member of the interval I and is such that $b - \epsilon < x$. Thus, to each $\epsilon > 0$ we have produced an $x \in I$ for which $b - \epsilon < x$. By virtue of Proposition A.5, we conclude that $b = \sup I$.

Of course, we have notion parallel to the supremum for sets which are bounded below.

Definition A.6. Let A be a set of real numbers which is bounded below. Suppose that there exists a number I with the following two properties:

- i. I is a lower bound of A .
- ii. For any lower bound N of A , $N \leq I$.

Then I is called the *infimum* (or *greatest lower bound*) of the set A and we write $I = \inf A$.

As with the definition of supremum, a set which is bounded below has at most one infimum and the justification of this parallels (almost exactly) the proof of Lemma A.3. The question of existence is dealt with by the following corollary to Theorem A.4 and whose proof I'll leave as an exercise.

Corollary A.7. Suppose A is a non-empty subset of \mathbb{R} which is bounded below. Then

$$I = \inf A$$

exists.

Exercise 15

Prove Corollary A.7 by completing the following steps.

1. Given a non-empty subset A which is bounded below, define the set

$$(-A) = \{x \in \mathbb{R} : x = -a \text{ for some } a \in A\}.$$

Prove that $-A$ is bounded above.

2. By making an appeal to Theorem A.4, let $S = \sup(-A)$. Prove that $-S = \inf A$ and so the infimum of A exists as the corollary asserts.

As we did for sets which were unbounded from above, if a non-empty set A is not bounded below, we will assume the standard convention and write

$$\inf A = -\infty.$$

Analogous to Proposition A.5, we have the following proposition for infima.

Proposition A.8. Let A be a non-empty set of real numbers which is bounded below and let I be a lower bound for A . Then $I = \inf A$ if and only if the following condition is satisfied. For each $\epsilon > 0$, there exists $x \in A$ for which $x < I + \epsilon$.

Exercise 16

Prove the proposition above. You may prove it directly (as I did for Proposition A.5) or you may prove it using the idea used to prove Corollary A.7, i.e., by considering the set $-A$.

We shall now discuss how suprema and infima are used to gain information about real-valued functions. To this end, let I be a non-empty set (often I will be a subset of the real line) and let $f : I \rightarrow \mathbb{R}$, i.e., f is a function mapping the set I into the set \mathbb{R} . We will say that f is *bounded above* if the set of real numbers

$$f(I) := \{f(x) : x \in I\}$$

is bounded above. Any upper bound for $f(I)$ will also be called an upper bound of the function f . In this case, we will write

$$\sup_{x \in I} f(x) = \sup f(I),$$

called the supremum of f on the interval I . We say that f is *bounded below* if the set $f(I)$ is bounded below. Any lower bound for $f(I)$ will also be called a lower bound of the function f . Here we write

$$\inf_{x \in I} f(x) = \inf f(I)$$

which is called the infimum of f on the interval I . Finally, we say that f is *bounded* if $f(I)$ is a bounded set.

Remark A.9. In the special case that I is the interval $[a, b]$, we will write

$$\sup_{a \leq x \leq b} f(x) = \sup_{x \in I} f(x) \quad \text{and} \quad \inf_{a \leq x \leq b} f(x) = \inf_{x \in I} f(x).$$

We will assume a similar notion when $I = (a, b)$, $[a, b)$ or (a, b) .

Exercise 17

Let $f : I \rightarrow \mathbb{R}$. Prove that f is bounded if and only if the absolute value of f , $x \mapsto |f(x)|$ is bounded above and further, in the case that either of these equivalent conditions are satisfied, prove that

$$\sup_{x \in I} |f(x)| = \max\{M_s, M_i\}$$

where

$$M_s = \left| \sup_{x \in I} f(x) \right| \quad \text{and} \quad M_i = \left| \inf_{x \in I} f(x) \right|.$$

In light of the above definition, a function f on I is bounded if and only if $\sup_{x \in I} |f(x)|$ exists as a real number. In this case, we write

$$\|f\|_\infty = \sup_{x \in I} |f(x)|$$

which is called the *sup-norm* of f . We will denote by $B(I)$ the set of bounded real-valued functions on the interval I . In the above notation, we can write

$$B(I) = \{f : I \rightarrow \mathbb{R} : \|f\|_\infty < \infty\}.$$

B The Riemann-Darboux integral

In this short section, we cover the basic properties of the Riemann integral⁵ needed for our study of Fourier analysis (at a level suited for the course textbook). As stated in lecture, it turns out that even the Riemann integral –the integral you’ve known and studied since your first brush with calculus – is insufficient for a comprehensive theory of Fourier analysis. To treat the comprehensive theory, in earnest, one needs the Lebesgue theory of integration. Though we will try to explore the necessity of Lebesgue integration later (while illustrating the shortcomings of the Riemann integral), we first need to lay the groundwork for the Riemann integral. This is the subject to which we now turn.

Consider an interval $I = [a, b]$ of \mathbb{R} . A *partition* P of I is a finite subset $P = \{x_0, x_1, x_2, \dots, x_N\}$ of I such that

$$a = x_0 < x_1 < x_2 < \dots < x_{N-1} < x_N = b.$$

⁵Technically, the integral we study is called the Riemann-Darboux integral, a name which also gives credit to the French mathematician Jean Gaston Darboux.

Though a partition P is simply a finite subset of I which is enumerated and includes the endpoints a and b , you should picture P as dividing up the interval I into the subintervals $[x_{n-1}, x_n]$ for $n = 1, 2, \dots, N$. Given a bounded function $f : I \rightarrow \mathbb{R}$ and a partition P of I , define

$$m_n = \inf_{x_{n-1} \leq x \leq x_n} f(x) \quad \text{and} \quad M_n = \sup_{x_{n-1} \leq x \leq x_n} f(x)$$

for each $n = 1, 2, \dots, N$. With these, we define the upper and lower (Darboux) sums of f with respect to the partition P respectively by

$$U(f, P) = \sum_{n=1}^N M_n(x_n - x_{n-1}) \quad \text{and} \quad L(f, P) = \sum_{n=1}^N m_n(x_n - x_{n-1}).$$

Of course, both $U(f, P)$ and $L(f, P)$ exist because f is a bounded function by virtue of the preceding exercise. The numbers $U(f, P)$ and $L(f, P)$ are respectively overestimates and underestimates for the area under the graph of f on the interval I , when this area is a sensible notion. These estimates are produced by forming rectangles above and below the graph of f where the width of the rectangles are determined by the subdivisions of I produced by the partition P . By properties of the supremum and infimum, observe that

$$L(f, P) \leq U(f, P) \tag{4}$$

and this holds for every partition P and every bounded function $f : I \rightarrow \mathbb{R}$.

Given partitions P and Q of I , Q is said to be a *refinement* of P if $P \subseteq Q$. As the partition P divides the interval I by subintervals, you should think of the refinement Q as producing finer subdivisions – hence the name “refinement”. With the aim of comparing upper and lower sums, we need the following lemma. The lemma says essentially that finer divisions of I yields “better” estimates for the area under the graph of f .

Lemma B.1. *Let P and Q be partitions of I and suppose that Q is a refinement of P . For any $f \in B(I)$,*

$$L(f, P) \leq L(f, Q) \quad \text{and} \quad U(f, Q) \leq U(f, P).$$

For brevity, I have omitted the proof of the preceding lemma. The idea for the proof is to start with a partition P and first consider a refinement Q_1 which contains exactly one more point than P . In this case, it is easy to establish the estimates of the lemma for $Q = Q_1$. Once this is shown, it’s just a matter of inductively adding points to Q_1 and “chaining” the resulting estimates together until the desired estimate is reached. The reader is referred to Theorem 6.4 of [?] for the proof.

Thinking back to our picture of the area under the graph, which we will soon interpret as the integral, we expect the lower sums to be underestimates for this area and the upper sums to be overestimates. Equivalently, we can start to think of the integral as a number which sits below all of the upper sums and above all of the lower sums. To think about how to approximate this number, we need to invoke the notion of supremum and infimum. To this end, we’ll need another lemma which will help us to make sure the infimum and supremum exist.

Lemma B.2. *Let $f \in B(I)$ and let P and Q be partitions of I . Then*

$$\left(\inf_{x \in I} f(x) \right) (b - a) \leq L(f, P) \leq U(f, Q) \leq \left(\sup_{x \in I} f(x) \right) (b - a)$$

Proof. We first note that the trivial partition $T = \{a, b\} = \{x_0, x_1\}$ has

$$L(f, T) = \sum_{n=1}^1 m_n(x_n - x_{n-1}) = m_1(x_1 - x_0) = \left(\inf_{x_0 \leq x \leq x_1} f(x) \right) (x_1 - x_0) = \left(\inf_{x \in I} f(x) \right) (b - a)$$

and

$$U(f, T) = \sum_{n=1}^1 m_n(x_n - x_{n-1}) = m_1(x_1 - x_0) = \left(\sup_{x_0 \leq x \leq x_1} f(x) \right) (x_1 - x_0) = \left(\sup_{x \in I} f(x) \right) (b - a).$$

Thus, for any partitions P and Q , Lemma B.1 guarantees that

$$\left(\inf_{x \in I} f(x) \right) (b - a) = L(f, T) \leq L(f, P)$$

and

$$U(f, Q) \leq \left(\sup_{x \in I} f(x) \right) (b - a)$$

because P and Q are necessarily refinements of T . It remains to establish the inner inequality.

To this end, observe that the union $R = P \cup Q$ is also a partition of I for it is necessarily a finite subset of I which contains a and b . Further, R is a refinement of both partitions P and Q . Thus, by another appeal to Lemma B.1 and in view of (4), we have

$$L(f, P) \leq L(f, R) \leq U(f, R) \leq U(f, Q)$$

which guarantees that $L(f, P) \leq U(f, Q)$ as was asserted. \square

Let's isolate some conclusions of the preceding lemma. First, it says that, for any partition P of I ,

$$L(f, P) \leq \left(\sup_{x \in I} f(x) \right) (b - a).$$

Hence, the set

$$\{L(f, P) : P \text{ is a partition of } I\}$$

is bounded above and, in view of Theorem A.4, its supremum exists. Thus, we define

$$\int_I f(x) dx = L(f) = \sup_P L(f, P)$$

where this supremum is taken over all partitions P of I . This is called the lower Darboux sum of f on I . Analogously, Lemma B.2 guarantees that the infimum of all upper sums exists and so we define the upper Darboux sum of f on I as

$$\overline{\int_I f(x) dx} = U(f) = \inf_P U(f, P)$$

As we've established quite a few inequalities involving upper and lower sums pertaining to the same and different partitions of I , it's helpful to have some sense of how $U(f)$ and $L(f)$ compare for a given bounded function $f : I \rightarrow \mathbb{R}$. To this end, let's fix a partition Q of I and note that, in view of Lemma B.2,

$$L(f, P) \leq U(f, Q)$$

for all partitions P of I . Thus, $U(f, Q)$ is an upper bound of the set of real numbers $\{L(f, P) : P \text{ is a partition of } I\}$. By the defining property of the supremum, we have

$$L(f) = \sup_P L(f, P) \leq U(f, Q).$$

Noting however that Q was arbitrary, we see that $L(f)$ is a lower bound for $U(f, Q)$ for all partitions Q of I . By the defining property of the infimum, we have

$$L(f) \leq \inf_Q U(f, Q) = U(f).$$

Let's summarize this information.

Proposition B.3. Let $f : I \rightarrow \mathbb{R}$ be a bounded function, i.e., $f \in B(I)$. Then the upper and lower Darboux sums,

$$\overline{\int_I} f(x) dx = U(f) = \inf_P U(f, P)$$

and

$$\underline{\int_I} f(x) dx = L(f) = \sup_P L(f, P),$$

exist. Furthermore,

$$\underline{\int_I} f(x) dx \leq \overline{\int_I} f(x) dx.$$

Exercise 18

This exercise will give you an idea of what's going on in the above construction. In what follows, we will focus on the interval $I = [0, 1]$. For each $N = 1, 2, \dots$, we shall consider the (regular) partition

$$P_N = \{x_0 < x_1 < \dots < x_N = 1\} = \left\{x_n = \frac{n}{N} : n = 0, 1, 2, \dots, N\right\}$$

of the interval I .

1. For the function $f(x) = 1$ for $0 \leq x \leq 1$, compute $U(f, P_N)$ and $L(f, P_N)$.

(a) Is it true that $L(f, P_N) \leq U(f, P_N)$?

(b) Show that $\lim_{N \rightarrow \infty} (U(f, P_N) - L(f, P_N)) = 0$.

2. For the function $f(x) = x$ for $0 \leq x \leq 1$, compute $U(f, P_N)$ and $L(f, P_N)$.

(a) Is it true that $L(f, P_N) \leq U(f, P_N)$?

(b) Show that $\lim_{N \rightarrow \infty} (U(f, P_N) - L(f, P_N)) = 0$.

3. For the Dirichlet function f defined by

$$f(x) = \begin{cases} 1 & x \in \mathbb{Q} \\ 0 & x \notin \mathbb{Q} \end{cases}$$

for $0 \leq x \leq 1$, compute $U(f, P_N)$ and $L(f, P_N)$.

(a) Is it true that $L(f, P_N) \leq U(f, P_N)$?

(b) Does $\lim_{N \rightarrow \infty} (U(f, P_N) - L(f, P_N)) = 0$?

4. For the first two examples above, you've seen a sequence (an enumerated collection) of partitions $\{P_n\}$ for which

$$\lim_{N \rightarrow \infty} (U(f, P_N) - L(f, P_N)) = 0.$$

In view of Proposition B.3 and the above fact, does it suffice to conclude that

$$\underline{\int_I} f(x) dx = \overline{\int_I} f(x) dx?$$

Prove your assertion (or find a counter example).

5. Is it true that if there is a sequence of partitions $\{P_N\}$ for which

$$\lim_{N \rightarrow \infty} (U(f, P_N) - L(f, P_N)) \neq 0,$$

then

$$\int_I f(x) dx \neq \overline{\int_I f(x) dx}?$$

Prove your assertion (or find a counter example).

Finding motivation in the preceding example and returning again to our intuition of areas, we would hope that a sensible notion of area under the graph could be gotten by approximating the area from above by upper sums and from below by lower sums. Thus, if such an area does exist, we would hope that the supremum of all the lower sums coincides with the supremum of all the lower sums and so the inequality of the preceding proposition is actually an equality. This is exactly the right idea and we give this situation a name.

Definition B.4. Let $f \in B(I)$ and let

$$U(f) = \overline{\int_I f(x) dx} \quad \text{and} \quad L(f) = \underline{\int_I f(x) dx}$$

be its upper and lower Darboux sums. We say that f is Riemann integrable on I and write $f \in R(I)$ if $U(f) = L(f)$. In this case, the Riemann integral of f is defined to be the number

$$\int_I f(x) dx = U(f) = L(f).$$

By virtue of Proposition A.5 and its analogue for infima, we have the following characterization for integrability.

Proposition B.5. Let $f \in B(I)$. Then $f \in R(I)$ (that is, Riemann integrable) if and only if the following conditions is satisfied:

For each $\epsilon > 0$, there is a partition P_ϵ of I for which $U(f, P_\epsilon) - L(f, P_\epsilon) < \epsilon$.

Proof. We first suppose that f is Riemann integrable. By Proposition A.5, let Q_1 be a partition for which $L(f) - L(f, Q_1) < \epsilon/2$. Similarly, by the characterization for infimum, let Q_2 be a partition of I for which $U(f, Q_2) - U(f) < \epsilon/2$. With these partitions in mind, we set $P_\epsilon = Q_1 \cup Q_2$ and observe that P_ϵ is a refinement of both Q_1 and Q_2 . By Lemma B.1, we have $L(f, P_\epsilon) \geq L(f, Q_1)$ and $U(f, P_\epsilon) \leq U(f, Q_2)$ and thus

$$U(f, P_\epsilon) - L(f, P_\epsilon) \leq U(f, Q_2) - L(f, Q_1) < U(f) + \epsilon - (L(f) - \epsilon/2) = U(f) - L(f) + \epsilon.$$

Of course, because $f \in R(I)$, $U(f) = L(f)$ and so the above inequality shows that $U(f, P_\epsilon) - L(f, P_\epsilon) < \epsilon$.

Conversely, let's assume that the desired property holds. Let $\epsilon > 0$, and using the property select a partition P for which $U(f, P) - L(f, P) < \epsilon$. As $U(f)$ and $L(f)$ are constructed from infima and suprema respectively, we have

$$U(f) - L(f) \leq U(f, P) - L(f, P) < \epsilon.$$

In view of Proposition B.3, we also have $U(f) - L(f) \geq 0$. Hence, to each $\epsilon > 0$, we have

$$0 \leq U(f) - L(f) < \epsilon.$$

We may therefore conclude that $U(f) = L(f)$ for the only number "lodged" between zero and every positive number is the number zero itself. \square

Let's now introduce the notions of integration and integrability for complex-valued functions.

Definition B.6. Let $I = [a, b]$ and consider a complex-valued function $f : I \rightarrow \mathbb{C}$. In this case f is necessarily of the form

$$f(x) = u(x) + iv(x)$$

for $x \in I$ where $u, v : I \rightarrow \mathbb{R}$. We saw that f is Riemann integrable on I if u and v are Riemann integrable on I and we define the integral of f on I to be the complex number

$$\int_I f(x) dx = \left(\int_I u(x) dx \right) + i \left(\int_I v(x) dx \right).$$

With a slight abuse of notation, we write $f \in R(I)$ and so $R(I)$ is then taken to be the set of Riemann-integrable complex-valued functions on I . We will also use the notations

$$\int_I f = \int_a^b f = \int_a^b f(x) dx$$

to denote the integral of f .

Let's make a few notes concerning the above definition. First, the functions u and v are called the real and imaginary parts of f respectively. We'll often write $f = \operatorname{Re}(f) + i \operatorname{Im}(f)$ where $\operatorname{Re}(f) = u$ and $\operatorname{Im}(f) = v$. In the (special) case in which f is a real-valued function from I to \mathbb{R} , we can write $f = \operatorname{Re}(f) + i \operatorname{Im}(f) = \operatorname{Re}(f) + i0 = f + i0$ and so here

$$\int_I f = \int \operatorname{Re}(f) + i \int_I 0 = \int_I \operatorname{Re}(f)(x) dx + i0 = \int_I \operatorname{Re}(f)(x) dx$$

because the integral of the zero function is just 0. In this way we observe that the definition of the Riemann integral for complex-valued functions is an extension of the Riemann integral for real-valued functions (as it recaptures the real-valued version of the Riemann integral).

Now that we know what integrability means, it's high time to give some properties of the integral.

Proposition B.7. Let $I = [a, b] \subseteq \mathbb{R}$.

1. For any complex numbers α and β and any $f, g \in R(I)$, the linear combination $\alpha f + \beta g \in R(I)$ and

$$\int_I (\alpha f + \beta g) = \alpha \int_I f + \beta \int_I g.$$

This says that $R(I)$ is a vector space over \mathbb{C} and the integral (viewed as a function $f \rightarrow \int_I f$) is linear map from $R(I)$ to \mathbb{C} .

2. If $f, g \in R(I)$, then the product $fg \in R(I)$.
3. Constant functions are Riemann-integrable and for any constant function $x \mapsto \alpha$ where $\alpha \in \mathbb{C}$,

$$\int_I \alpha = \alpha(b - a).$$

4. The set of continuous functions $C(I)$ are Riemann integrable. That is, $C(I) \subseteq R(I)$.

Proof. As the first statement was partially covered in Homework 1 (see also the exercise below), I'll omit the proof and refer the reader to [?] for details. See [?] for the proof of Item 2, as well. I will prove Items 3 and 4 here.

3. Let's first consider the constant function 1. This function is obviously bounded and, as it is real-valued, let's show that it is integrable by computing its upper and lower sums. For any partition $P = \{a = x_0 < x_1 < x_2 < \dots < x_N = b\}$, we have

$$M_n = \sup_{x_{n-1} \leq x \leq x_n} 1 = 1 = \inf_{x_{n-1} \leq x \leq x_n} 1 = m_n$$

for each $n = 1, 2, \dots, n$ and therefore

$$U(1, P) = \sum_{n=1}^N M_k(x_n - x_{n-1}) = \sum_{n=1}^N 1(x_n - x_{n-1}) = b - a$$

and

$$L(1, P) = \sum_{n=1}^N m_n(x_n - x_{n-1}) = \sum_{n=1}^N 1(x_n - x_{n-1}) = b - a.$$

Since the above is true for any partition P , we have

$$U(1) = \inf_P U(1, P) = \inf_P (b - a) = b - a = \sup_P (b - a) = \sup_P L(1, P) = L(f)$$

from which we conclude that the constant function 1 is Riemann-integrable and

$$\int_I 1 = b - a.$$

Now, given any complex number α , $\alpha = \alpha \cdot 1$ and so, by Item 1 and the fact that $1 \in R(I)$, $\alpha \in R(I)$ and

$$\int_I \alpha = \int_I \alpha \cdot 1 = \alpha \int_I 1 = \alpha(b - a)$$

as desired.

4. We will begin by proving the result for continuous real-valued functions. To this end, let $g : I \rightarrow \mathbb{R}$ be continuous on the interval $I = [a, b]$. Our proof makes use of two essential results from analysis, both of which rely on the interval I being closed and bounded (compact). First, in view of Theorem 4.15 of [?], g is necessarily a bounded function. Second, by virtue of Theorem 4.19 of [?], g is uniformly continuous on I . That is, to each positive number $\epsilon > 0$, there is $\delta > 0$ for which

$$|g(x) - g(y)| < \epsilon \quad \text{whenever} \quad |x - y| < \delta.$$

With these facts in mind, let's show that g is Riemann-integrable by meeting the equivalent condition of Proposition B.5. To this end, we note that g is bounded and we fix $\epsilon > 0$. In view of the uniform continuity of g , let $\delta > 0$ be such that

$$|g(x) - g(y)| < \frac{\epsilon}{2(b-a)} \quad \text{whenever} \quad |x - y| < \delta.$$

With this δ , let's consider the "regular" partition

$$P = \{a = x_0 < x_1 < \dots < x_N = b\} = \left\{ a + \frac{n}{N}(b-a) : n = 0, 1, \dots, N \right\}$$

where $N \in \mathbb{N}$ is chosen so that $N > (b-a)/\delta$ and so $(b-a)/N < \delta$. By this choice, let's make some observations. First, for any $n = 1, 2, \dots, N$, if

$$x, y \in [x_{n-1}, x_n] = \left[a + \frac{n-1}{N}(b-a), a + \frac{n}{N}(b-a) \right], \quad \text{then} \quad |x - y| < \frac{(b-a)}{N} < \delta.$$

and so $|g(x) - g(y)| < \epsilon/2(b-a)$ in view of the uniform continuity of g . So, for each $n = 1, 2, \dots, N$ and $y \in [x_{n-1}, x_n]$,

$$g(y) - \frac{\epsilon}{2(b-a)} < g(x) < \frac{\epsilon}{2(b-a)} + g(y)$$

and so $\epsilon/2(b - a) + g(y)$ is an upper bound for $\{g(x) : x \in [x_{n-1}, x_n]\}$. Consequently,

$$M_n = \sup_{x_{n-1} \leq x \leq x_n} g(x) \leq \frac{\epsilon}{2(b - a)} + g(y)$$

in view of the definition of the supremum. Because $y \in [x_{n-1}, x_n]$ was arbitrary, the above inequality shows that $M_n - \epsilon/2(b - a)$ is a lower bound for the set $\{g(y) : y \in [x_{n-1}, x_n]\}$ and so

$$M_n - \frac{\epsilon}{b - a} \leq \inf_{x_{n-1} \leq y \leq x_n} g(y) = m_n.$$

In this way we have established that

$$M_n - m_n < \frac{\epsilon}{2(b - a)}$$

for each $n = 1, 2, \dots, N$. Correspondingly,

$$\begin{aligned} U(g, P) - L(g, P) &= \sum_{n=1}^N M_k(x_n - x_{n-1}) - \sum_{n=1}^N m_n(x_n - x_{n-1}) \\ &= \sum_{n=1}^N (M_n - m_n)(x_n - x_{n-1}) \\ &< \sum_{n=1}^N \frac{\epsilon}{2(b - a)}(x_n - x_{n-1}) = \frac{\epsilon}{2(b - a)} \sum_{n=1}^N (x_n - x_{n-1}) = \frac{\epsilon}{2} < \epsilon. \end{aligned}$$

In view of Proposition B.5, we can therefore conclude that $g \in R(I)$.

Finally, let $f \in C(I)$ be a arbitrary complex-valued continuous function on I . An appeal to your result from Homework 1 shows that the real and imaginary parts of f , $\text{Re}(f)$ and $\text{Im}(f)$ are necessarily continuous real-valued functions on I . Thus, by virtue of our result in the previous paragraph, $\text{Re}(f)$ and $\text{Im}(f)$ are both Riemann-integrable. By definition (of integrability for complex-valued functions), we conclude that $f \in R(I)$. □

Exercise 19

In this exercise, you prove the real-valued analogue of the scalar multiplication portion of Item 1 of the proposition above. Throughout this exercise, c is a real number.

1. First, given a non-empty bounded set A of \mathbb{R} , we denote by cA the set of numbers of the form $c \cdot a$ where $a \in A$. That is, $cA = \{x \in \mathbb{R} : x = ca \text{ for } a \in A\}$. If $c > 0$, prove that

$$\sup cA = c \sup A \quad \text{and} \quad \inf cA = c \inf A.$$

2. If $c < 0$, formulate and prove an analogous statement for $\sup cA$ and $\inf cA$.
3. For the remainder of this exercise, $g : I \rightarrow \mathbb{R}$ will be an arbitrary bounded function. We will assume now that $c > 0$ and denote by cg the real-valued function on I defined by $(cg)(x) = cg(x)$ for $x \in I$. Use your result from Item 1 to prove that

$$U(cg, P) = cU(g, P) \quad \text{and} \quad L(cg, P) = cL(g, P).$$

for any partition P of I .

4. Continuing under the assumption that $c > 0$, prove that $U(cg) = c \cdot U(g)$ and $L(cg) = c \cdot L(g)$.

5. Use the item above to prove that, if $c > 0$, $g \in R(I)$ if and only if $cg \in R(I)$ and

$$c \int_I g = \int_I cg.$$

6. Comment on how the previous steps change if we allow c to be non-positive. In particular, is it still true that $cg \in R(I)$ if and only if $g \in R(I)$?

Another important property of the integral is captured by the following proposition.

Proposition B.8. *Let $f \in R(I)$, then the function $|f| : I \rightarrow \mathbb{R}$ defined by*

$$|f|(x) = |f(x)| = \sqrt{(\operatorname{Re}(f(x)))^2 + (\operatorname{Im}(f(x)))^2} \quad \text{for } x \in I$$

is Riemann integrable and

$$\left| \int_I f \right| \leq \int_I |f|.$$

As it is somewhat involved, we will not show that $f \in R(I)$ guarantees that $|f| \in R(I)$. For this, we refer the reader to Theorem 6.13 of [?]. We will however prove the inequality. We first need a lemma.

Lemma B.9. *Let $h_1, h_2 \in R(I)$ be real-valued functions such that $h_1(x) \leq h_2(x)$ for all $x \in I$. Then*

$$\int_I h_1 \leq \int_I h_2.$$

Exercise 20

Prove the lemma above. Hint: Start by showing that non-negative functions have non-negative integrals. Then use Item 1 of Proposition B.7.

Proof. Let f be an arbitrary complex-valued Riemann-integrable function on I and, in accordance with the remark preceding Lemma B.9 we will take for granted that $|f| \in R(I)$. In view of Exercise B (below), there is $\theta \in (-\pi, \pi)$ for which

$$\left| \int_I f \right| = e^{-i\theta} \left(\int_I f \right).$$

In view of Item 1 of Proposition B.7, this guarantees that

$$\left| \int_I f \right| = \int_I e^{-i\theta} f = \int_I (e^{-i\theta} f(x)) \, dx = \int_I \operatorname{Re}(e^{-i\theta} f(x)) \, dx + i \int_I \operatorname{Im}(e^{-i\theta} f(x)) \, dx.$$

As the left hand side of the above equation is purely real, this ensures that the purely imaginary part of the right hand side is zero and therefore

$$\left| \int_I f \right| = \int_I \operatorname{Re}(e^{-i\theta} f(x)) \, dx.$$

Now, for each $x \in I$,

$$\operatorname{Re}(e^{-i\theta} f(x)) \leq \sqrt{(\operatorname{Re}(e^{i\theta} f(x)))^2 + (\operatorname{Im}(e^{-i\theta} f(x)))^2} = |e^{-i\theta} f(x)| = |f(x)|$$

where we have used the fact that $|zw| = |z||w|$ for complex numbers z, w . Thus, by Lemma B.9, we have

$$\left| \int_I f \right| \leq \int_I \operatorname{Re}(e^{-i\theta} f(x)) \, dx \leq \int_I |f(x)| \, dx = \int_I |f|$$

as desired. □

Exercise 21

Prove that, for each complex number $z = a + ib \in \mathbb{C}$, there exists $\theta \in (-\pi, \pi]$ for which

$$e^{-i\theta} z = |z| = \sqrt{a^2 + b^2}.$$

In this way, every complex-number z can be written as

$$z = |z|e^{i\theta}$$

for some $\theta \in (-\pi, \pi]$ called the *phase*^a of z .

^aWhen $z \neq 0$, θ can be shown to be unique in this range.

Our next proposition is often called the “change of variables formula”. Because the proof is somewhat technical (and is actually best done in the context of the Riemann-Steiltjes integral), I have decided to omit it.

Proposition B.10 (Change of variables formula, Theorem 6.19 of [?]). *Let $A < B$ and $a < b$ be real numbers and suppose that $h : [A, B] \rightarrow [a, b]$ is a strictly increasing function mapping $[A, B]$ onto $[a, b]$ with derivative $h' \in R([A, B])$. Also, let $f \in R([a, b])$. Then the function $x \mapsto (f \circ h)(x)h'(x) = f(h(x))h'(x)$ is integrable on $[A, B]$ and*

$$\int_a^b f(x) dx = \int_{[a,b]} f = \int_{[A,B]} (f \circ h) \cdot h' = \int_A^B f(h(x))h'(x) dx$$

It should be noted that the proposition above has a very beautiful generalization to integration in \mathbb{R}^d in which the derivative h' is replaced by the Jacobean determinant of h 's d -dimensional analogous. This generalization is an essential tool used in the theory of integration on manifolds.

We end this section by treating a nice result which says that each Riemann-integrable function is “close” in a certain sense to a continuous function.

Proposition B.11. *Suppose that $f \in R(I)$ and f is bounded by B , i.e., $|f(x)| \leq B$ for all $x \in I$. Then there exists a sequence of continuous functions $\{f_k\} \subseteq C(I)$ such that*

$$\sup_{x \in I} |f_k(x)| \leq B$$

for all $k \in I$ and

$$\lim_{k \rightarrow \infty} \int_I |f(x) - f_k(x)| dx = 0.$$

Proof. See Lemma 1.5 of Stein-Shakarchi appendix □

C The essence of convergence

In introductory calculus (Math 121/161 and 122/162), you learned about the notion of convergence for sequences of real numbers. This notion was captured by saying, given a sequence of real numbers $\{a_n\}$ and another real number a , the sequence $\{a_n\}$ converges to a if the terms of the sequence a_n can be made arbitrarily close to a by taking n sufficiently large. This idea is essentially unchanged when we talk about convergence of sequences of complex numbers. This is captured in the following definition.

Definition C.1. *Let $\{w_n\}$ be a sequence of complex numbers (written $\{w_n\} \subseteq \mathbb{C}$) and let w be another complex number. We say that the sequence $\{w_n\}$ converges to w if the following condition is satisfied. For all $\epsilon > 0$, there exists a natural number N (written $N \in \mathbb{N}$) for which*

$$|w_n - w| < \epsilon \quad \text{whenever} \quad n \geq N.$$

The essential difference between the definitions of convergence for real and complex numbers is the way that distance (and closeness) is measured. In the above definition, the symbol $|\cdot|$ means the complex modulus and is defined by

$$|z| = \sqrt{a^2 + b^2}.$$

for a complex number $z = a + ib$ where this symbol is taken to mean the absolute value when applied to real numbers. As you have already explored this in Homework 1, I won't expound upon convergence of real and complex numbers further here. Let's instead move into a discussion concerning convergence of functions, which is the main notion of interest for the discussion of Fourier series.

Just as we think of a sequence of complex numbers converging to another complex number, in studying convergence of functions, we are interested in the study of a sequence of complex-valued functions $\{f_n\}$ defined on some set I getting "close" to another function f . A moment's thought about this invokes many questions, primarily the question of what it means to be "close". To that end, we will examine several inequivalent notions of closeness and convergence for functions. The first of which (and one of the weakest) is captured by the following definition.

Definition C.2. *Let I be an interval of the real line and let $\{f_n\}$ be a sequence of complex-valued functions on I , i.e., $f_n : I \rightarrow \mathbb{C}$ for each $n = 1, 2, \dots$. Let $f : I \rightarrow \mathbb{C}$ be another function. We say that the sequence $\{f_n\}$ converges to f pointwise on I if, for each $x \in I$,*

$$\lim_{n \rightarrow \infty} f_n(x) = f(x).$$

The important thing to note about the above definition is that the x is chosen before the limit is taken. Stated with ϵ 's and N 's, the above definition is as follows:

The sequence of functions f_n converges to f pointwise on I if, for each $\epsilon > 0$ and $x \in I$, there is an $N \in \mathbb{N}$ (depending on both ϵ and x) for which

$$|f_n(x) - f(x)| < \epsilon \quad \text{whenever} \quad n \geq N.$$

Example 1

In this example, we consider a sequence of real-valued functions converging pointwise on the interval $I = [0, 1]$. For each natural number n , define $f_n : I \rightarrow \mathbb{R} \subseteq \mathbb{C}$

$$f_n(x) = x^n$$

for $x \in I$ and $n \in \mathbb{N}$. We observe that, for $0 \leq x < 1$,

$$\lim_{n \rightarrow \infty} f_n(x) = \lim_{n \rightarrow \infty} x^n = 0$$

and, for $x = 1$,

$$\lim_{n \rightarrow \infty} f_n(x) = \lim_{n \rightarrow \infty} 1^n = 1.$$

Thus, our sequence of functions converges uniformly to the function $f : I \rightarrow \mathbb{R}$ defined by

$$f(x) = \begin{cases} 0 & 0 \leq x < 1 \\ 1 & x = 1 \end{cases}$$

for $x \in I$. The graphs of f_n are illustrated for $n = 1, 2, \dots, 20$ in Figure 9.




Figure 9: A famous picture: The graphs of $f_n(x) = x^n$ for $n = 1, 2, \dots, 20$.

It is important to note that each function f_n is continuous on I , however, the limit function f is not continuous on I . This illustrates that nice properties like continuity can be “broken” under taking pointwise limits.

A much stronger notion of convergence is captured by the following definition.

Definition C.3. Let $\{f_n\}$ be a sequence of complex-valued functions on I . Let $f : I \rightarrow \mathbb{C}$ be another complex-valued function on I . We say that the sequence $\{f_n\}$ converges uniformly to f on I if, for all $\epsilon > 0$ there exists $N \in \mathbb{N}$ for which

$$|f_n(x) - f(x)| < \epsilon \quad \text{whenever} \quad x \in I \text{ and } n \geq N.$$

In contrast to the definition of pointwise convergence, the definition of convergence requires that the integer N depend only on ϵ and be independent of $x \in I$. This notion is illustrated in Figure 10. In the figure, we see the graph of a real-valued function f (in black) in the center of a “band” of radius ϵ (in red). For a sequence of functions $\{f_n\}$ to converge uniformly to f (on an interval) means that, for sufficiently large n , the graph of f_n is completely contained in the band of radius ϵ surrounding f ; the blue line is an example of the graph of one such f_n .

We further illustrate this definition with some examples.

Example 2

Consider the sequence $\{f_n\}$ of functions defined on the interval $I = [-\pi, \pi]$ by

$$f_n(x) = \cos(x/n) - 1/2$$

for $x \in I$ and $n \in \mathbb{N}$. The graphs of f_n are illustrated for $n = 1, 2, \dots, 10$ in Figure 11.




Figure 10: An illustration of uniform convergence




Figure 11: The graphs of $f_n(x) = \cos(x/n) - 1/2$ for $n = 1, 2, \dots, 10$.

The figure suggests that the sequence $\{f_n\}$ converges to the constant function $f(x) = 1/2$ as $n \rightarrow \infty$. Let's prove that, not only does it converge to $f(x) = 1/2$, it does so uniformly.

Let $\epsilon > 0$ and select $N \in \mathbb{N}$ such that $N > \pi/\sqrt{\epsilon}$. Recalling the inequality for cosine,

$$|\cos(\theta) - 1| \leq |\theta^2| \quad \text{for all } \theta \in \mathbb{R}$$

which can be gotten from the mean value theorem or the racetrack principle, we observe that, for any $n \geq N$ and $x \in I = [-\pi, \pi]$,

$$|f_n(x) - f(x)| = |\cos(x/n) - 1/2 - 1/2| = |\cos(x/n) - 1| \leq \frac{x^2}{n^2} \leq \frac{\pi^2}{n^2} < \epsilon$$

because $n^2 \geq N^2 > \pi^2/\epsilon$. The careful reader should note that the above estimate holds for all $x \in I$ and for all $n \geq N$ (and not for a particular x). We have shown that the sequence $\{f_n\}$ converges uniformly to $f(x) = 1/2$.

Exercise 22

Given an interval I , we recall the supremum norm defined, for $f : I \rightarrow \mathbb{C}$ by

$$\|f\|_\infty = \sup_{x \in I} |f(x)|.$$

In this exercise, you will prove that $\|\cdot\|_\infty$ is a *bona fide* norm on the space of bounded complex-valued functions on I .

1. Prove that, for any pair of bounded functions f and g ,

$$\|f + g\|_\infty \leq \|f\|_\infty + \|g\|_\infty.$$

2. Prove that, for each complex number α and bounded function $f : I \rightarrow \mathbb{C}$,

$$\|\alpha f\|_\infty = |\alpha| \|f\|_\infty$$

where $|\alpha|$ is the complex modulus of α .

3. Prove that, for a bounded function f , $\|f\|_\infty = 0$ if and only if $f(x) = 0$ for all $x \in I$.
4. Given a sequence $\{f_n\}$ of bounded complex-valued functions on I and $f : I \rightarrow \mathbb{C}$, prove that the sequence $\{f_n\}$ converges uniformly to f if and only if

$$\lim_{n \rightarrow \infty} \|f_n - f\|_\infty = 0.$$

As the notion of “Cauchy sequence” is essential for the convergence for complex-numbers and, in fact, provides a characterization for convergence as you proved in Homework 1, we have a similar Cauchy property for functions which characterizes uniform convergence. This characterization is outlined in the following theorem.

Theorem C.4. *Let $\{f_n\}$ be a sequence of complex-valued functions on an interval $I \subseteq \mathbb{R}$. The sequence $\{f_n\}$ converges uniformly (to some function f) on I if and only if it satisfies the following property:*

(UC) *For all $\epsilon > 0$, there exists a natural number N such that*

$$|f_n(x) - f_m(x)| < \epsilon \quad \text{whenever } x \in I \text{ and } n, m \geq N.$$

The equivalent property (UC) is called the Uniform Cauchy condition. Any sequence of functions $\{f_n\}$ satisfying the condition is said to be uniformly Cauchy on I .

Proof. Let us first assume that $\{f_n\}$ converges uniformly to a function f on I . Let $\epsilon > 0$ and by our supposition let N be a natural number for which

$$|f_n(x) - f(x)| < \epsilon/2$$

for all $n \geq N$ and $x \in I$. Then, for any $n, m \geq N$, we have

$$|f_n(x) - f_m(x)| = |f_n(x) - f(x) + f(x) - f_m(x)| \leq |f_n(x) - f(x)| + |f(x) - f_m(x)| \leq \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$$

for all $x \in I$. Thus the sequence $\{f_n\}$ is uniformly Cauchy on I .

Conversely, let's assume that the sequence $f_n(x)$ is uniformly Cauchy on I . This implies, in particular, that $\{f_n(x)\}$ is a Cauchy sequence of complex numbers for each $x \in I$. Because all Cauchy sequences of complex numbers converge, for each $x \in I$, the limit $\lim_{n \rightarrow \infty} f_n(x)$ exists and we will denote its value by $f(x)$, which is just a complex number. In this way, we produce a function $f : I \rightarrow \mathbb{C}$ simply by identifying each x with the value of the limit $\lim_{n \rightarrow \infty} f_n(x)$, i.e., defining

$$f(x) = \lim_{n \rightarrow \infty} f_n(x)$$

for each $x \in I$. So now we have a candidate (f) for the uniform limit. It remains to show that our sequence, in fact, converges uniformly to this f . To see this, we let $\epsilon > 0$ and choose a natural number N for which

$$|f_n(x) - f_m(x)| < \frac{\epsilon}{2}$$

for all $n, m \geq N$ and $x \in I$. Now, let $x \in I$ and $n \geq N$ be arbitrary (but fixed). The convergence of the numerical sequence $\{f_n(x)\}$ guarantees that there is some natural number $N_x \geq N$ for which

$$|f_m(x) - f(x)| < \frac{\epsilon}{2}$$

whenever $m \geq N_x$. In particular, this works when $m = N_x \geq N$ and so

$$|f_n(x) - f(x)| = |f_n(x) - f_{N_x}(x) + f_{N_x}(x) - f(x)| \leq |f_n(x) - f_{N_x}(x)| + |f_{N_x}(x) - f(x)| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$$

Thus, to each $\epsilon > 0$, we have found a natural number N for which

$$|f_n(x) - f(x)| < \epsilon$$

whenever $x \in I$ and $n \geq N$. Therefore, $\{f_n\}$ converges uniformly on I (to f). \square

The above theorem is extremely useful when one has a sequence of nice functions (which is uniformly Cauchy) but has no obvious candidate for the uniform limit. Here, of course, infinite series comes to mind.

Definition C.5. Let $\{f_n\}$ be a sequence of complex-valued functions on I . The (formal) sum $\sum_n f_n$ is called a series of functions. To investigate the convergence of $\sum_n f_n$, we define, for each $N = 1, 2, \dots$,

$$S_N(x) = \sum_{n=1}^N f_n(x) \quad \text{for } x \in I.$$

The functions S_1, S_2, \dots , form a sequence of complex-valued functions on I , $\{S_N\}$, called the sequence of partial sums for the series $\sum_n f_n$. If, for each $x \in I$, the limit

$$\lim_{N \rightarrow \infty} S_N(x)$$

exists, we say that the series $\sum_n f_n$ converges on I . In this case, the limit is a function $S : I \rightarrow \mathbb{R}$ defined by

$$S(x) = \lim_{N \rightarrow \infty} S_N(x) = \lim_{N \rightarrow \infty} \sum_{n=1}^N f_n(x)$$

and we write

$$\sum_{n=1}^{\infty} f_n(x) = S(x)$$

to denote this function, called the sum of the series. We say that the series $\sum_n f_n$ converges uniformly on I if its sequence of partial sums $\{S_N\}$ converges uniformly on I to the sum of the series.

As with numerical series, one can often learn that a series converges without ever knowing its sum. For instance, the integral test from calculus shows that the series of numbers

$$\sum_{n=1}^{\infty} \frac{1}{n^3}$$

converges (this is p -series for $p = 3$). Though it can be approximated to any degree of accuracy, its sum is unknown. With this in mind, it is important to have various tests for series (uniform) convergence without knowing the limit. The following corollary of Theorem C.4 gives us exactly this.

Corollary C.6 (Uniform Cauchy Criterion). *Let $\{f_n\}$ be a sequence of complex-valued functions on I and consider the series $\sum_n f_n$. The series $\sum_n f_n$ converges uniformly on I if and only if the following property is satisfied.*

For all $\epsilon > 0$ there is a natural number N for which

$$\left| \sum_{k=n}^{k=m} f_k(x) \right| < \epsilon$$

for all $x \in I$ and $m \geq n \geq N$. This property is called the Uniform Cauchy Criterion for the series $\sum_n f_n$.

Exercise 23

In this exercise, you will prove Corollary C.6 and then use the corollary to establish sufficient conditions for the absolute convergence of power series – things you will remember from calculus (M122).

- Using Theorem C.4, prove Corollary C.6.
- If a series $\sum_n f_n$ of functions $\{f_m\}$ converges uniformly on I , prove that $\{f_n\}$ converges uniformly to the zero function on I .
- For the remainder of this exercise, we fix a positive constant M and define $I = [-M, M] \subseteq \mathbb{R}$. Given a sequence of complex-numbers $\{c_n\}$, consider the sequence of complex-valued functions $\{f_n\}$ on I defined by

$$f_n(x) = \frac{c_n}{n!} x^n$$

for $x \in I$. If the sequence $\{c_n\}$ is bounded, i.e., $\sup_{n \in \mathbb{N}} |c_n| < \infty$, use Corollary C.6 (and no other convergence test) to prove that the series

$$\sum_{n=1}^{\infty} \frac{c_n}{n!} x^n$$

converges uniformly on I .

- Let $f : I \rightarrow \mathbb{C}$ be infinitely differentiable and assume that $\sup_{n=0,1,\dots} |f^{(n)}(0)| < \infty$; here $f^{(n)}(0)$ is the n^{th} -derivative of f at 0. Use the previous item to prove that the series

$$\sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n$$

converges uniformly on I . This series is called the Maclaurin series for f . (Your proof here should be approximately one sentence).

5. Looking back at Item 3, find a condition on the sequence $\{c_k\}$ which is less restrictive than boundedness and which still guarantees that the series

$$\sum_{n=1}^{\infty} \frac{c_n}{n!} x^n$$

converges uniformly on I . Hint: You should take a look at Stirling's formula (which you can take for granted as long as you interpret the formula/approximation correctly). If you're interested, a nice proof of Stirling's formula can be found in Exercise 5 of Homework 2 for my Math 122 class.

C.1 Properties of Uniform Convergence

In this short subsection, we discuss some properties preserved under uniform convergence. Specifically, we focus on continuity and integration. Let's consider a couple of examples.

Example 3

Given $0 < \delta < 1$, let $I_\delta = [-1 + \delta, 1 - \delta]$ and consider the series

$$\sum_{n=0}^{\infty} x^n$$

for $x \in I_\delta$. We claim that this series converges uniformly on I_δ to the function

$$f(x) = \frac{1}{1-x}. \quad (5)$$

To see this, we first observe that the partial sums $\{S_N\}$ satisfy the formula

$$S_N(x) = \sum_{n=0}^N x^n = \frac{1-x^{N+1}}{1-x}$$

for $x \in I_\delta$. The validity of this formula can be seen by multiplying both sides by $1-x$ and simplifying. To see that this series converges uniformly, let $\epsilon > 0$ and choose M to be a natural number for which $M > \ln(\epsilon\delta)/\ln(1-\delta)$. For any $x \in I_\delta$ and $N \geq M$, observe that

$$|f(x) - S_N(x)| = \left| \frac{1}{1-x} - \frac{1-x^{N+1}}{1-x} \right| = \frac{|x|^{N+1}}{|1-x|} \leq \frac{(1-\delta)^{N+1}}{\delta} < \epsilon$$

where we have used the fact that $N+1 > M \geq \ln(\epsilon\delta)/\ln(1-\delta)$. Therefore, we have proved that this series converges uniformly to f . I encourage you to show that this series converges uniformly using only Corollary C.6 (and not making reference to f).

An important thing to note about the above example is that, each $S_N(x)$ is continuous and the limit function $f(x) = 1/(1-x)$ is also continuous on the interval I_δ , a fact that was also true in the preceding example. This stands in contrast to the Example C in which the limit function failed to be continuous. As it turns out, this is a key difference between pointwise convergence and uniform convergence. This is detailed in the following theorem, whose proof can be found in [?] (see Theorem 7.12 therein).

Theorem C.7. *Let $\{f_n\}$ be a sequence of complex-valued functions on I and suppose that $\{f_n\}$ converges uniformly to a function $f : I \rightarrow \mathbb{C}$. If each function f_n is continuous, i.e., $\{f_n\} \subseteq C(I)$, then f is necessarily a continuous function.*

Let's explore some other important properties of uniform convergence. Our next result shows that uniform convergence plays nicely with the Riemann-Darboux integral.

Theorem C.8. *Let $\{f_n\}$ be a sequence of complex-valued functions which converges uniformly to a function $f : I \rightarrow \mathbb{C}$; here, $I = [a, b]$. If each function f_n is Riemann-integrable, i.e., $\{f_n\} \subseteq R(I)$, then f is Riemann-integrable and*

$$\lim_{n \rightarrow \infty} \int_I |f_n - f| = 0.$$

Further

$$\lim_{n \rightarrow \infty} \int_I f_n = \int f.$$

Proof. We first show that the limit f is Riemann-integrable by showing its real and imaginary parts, u and v are Riemann-integrable. For each n , denote by u_n and v_n the real and imaginary parts of f_n respectively. We will show that u and v are Riemann integrable by appealing to the $\epsilon - P$ characterization, Proposition B.5. Let's first focus on the real parts $\{u_n\}$ and u . Let $\epsilon > 0$ and, by the uniform convergence of $\{f_n\}$, let N be a natural number for which

$$|u_n(x) - u(x)| \leq \sqrt{(u_n(x) - u(x))^2 + (v_n(x) - v(x))^2} = |f_n(x) - f(x)| < \epsilon/4(b - a)$$

for all $x \in I$ and $n \geq N$. In particular, upon setting $u_0 = u_N$, this yields the inequality

$$u_0(x) - \frac{\epsilon}{4(b - a)} < u(x) < u_0(x) + \frac{\epsilon}{4(b - a)} \tag{6}$$

for all $x \in I$. This inequality implies that u is bounded on the interval I in view of our hypothesis that $u_0 = u_N \in R(I)$. By virtue of Proposition B.5, let P be a partition of I for which $U(u_0, P) - L(u_0, P) < \epsilon/2$. For this partition, the inequality (6) guarantees that

$$\begin{aligned} U(u, P) &= \sum_n \left(\sup_{x_{n-1} \leq x \leq x_n} u(x) \right) (x_n - x_{n-1}) \\ &\leq \sum_n \left(\sup_{x_{n-1} \leq x \leq x_n} u_0(x) + \frac{\epsilon}{4(b - a)} \right) (x_n - x_{n-1}) \\ &\leq U(u_0, P) + \sum_n \frac{\epsilon}{4(b - a)} (x_n - x_{n-1}) \\ &\leq U(u_0, P) + \frac{\epsilon}{4}. \end{aligned}$$

Similarly, the inequality (6) guarantees the analogous lower estimate

$$L(u_0, P) - \frac{\epsilon}{4} \leq L(u, P)$$

Together, these estimates guarantees that

$$U(u, P) - L(u, P) \leq U(u_0, P) - L(u_0, P) + \frac{\epsilon}{2} < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$$

and from this we can conclude that $u \in R(I)$. A completely analogous argument shows that $v \in R(I)$ and so, by the definition of Riemann-integrability for complex-valued functions, the limit function $f \in R(I)$.

Let us now prove the statements concerning the limit $\lim_{n \rightarrow \infty} \int_I |f_n - f|$. In view of the definition of the L^∞ -norm, we have

$$|f_n(x) - f(x)| \leq \|f_n - f\|_\infty$$

for all $x \in I$ and $n \in \mathbb{N}$. In view of Lemma B.9, we have

$$0 \leq \int_I |f_n(x) - f(x)| dx \leq \int_I \|f_n - f\|_\infty dx = (b - a)\|f_n - f\|_\infty.$$

Thus, by virtue of Exercise 9 and the squeeze theorem, the preceding inequality shows that

$$\lim_{n \rightarrow \infty} \int_I |f_n - f| = 0$$

because $\|f_n - f\|_\infty \rightarrow 0$ as $n \rightarrow \infty$.

Finally, by virtue of Propositions B.7 and B.8, we have

$$\left| \int_I f_n - \int_I f \right| = \left| \int_I (f_n - f) \right| \leq \int_I |f_n - f|$$

for all n . Another appeal to the squeeze theorem (and the preceding limit) guarantees that

$$\lim_{n \rightarrow \infty} \int_I f_n = \int_I f.$$

□

Corollary C.9. *Let $\{f_n\}$ be a sequence of complex-valued functions on $I = [a, b]$ and suppose that the series $\sum_{n=0}^\infty f_n$ converges uniformly on I . If each f_n is Riemann-integrable, then the sum of the series is Riemann-integrable and*

$$\int_I \sum_{n=0}^\infty f_n = \sum_{n=0}^\infty \int_I f_n.$$

Proof. The hypothesis that $\sum_{n=0}^\infty f_n$ converges uniformly means that the sequence of partial sums $\{S_N\}$ defined by

$$S_N(x) = \sum_{n=0}^N f_n(x)$$

for $x \in I$ converges uniformly on I . Also, the supposition that each f_n is Riemann-integrable guarantees that each partial sum is Riemann-integrable in view of Proposition B.7. By the (finite) linearity of the integral, we have

$$\int_I S_N = \sum_{n=0}^N \int_I f_n$$

for each natural number N . Thus, an appeal to the preceding theorem guarantees that the limit $\sum_{n=0}^\infty f_n = \lim_{N \rightarrow \infty} S_N$ is Riemann-integrable and

$$\int_I \sum_{n=0}^\infty f_n = \int_I \lim_{N \rightarrow \infty} S_N = \lim_{N \rightarrow \infty} \int_I S_N = \lim_{N \rightarrow \infty} \sum_{n=0}^N \int_I f_n;$$

in particular, the limit on the right exists. Of course, this is what it means for the series of the numbers $\int_I f_n$ to converge and so we have

$$\int_I \sum_{n=0}^\infty f_n = \lim_{N \rightarrow \infty} \sum_{n=0}^N \int_I f_n = \sum_{n=0}^\infty \int_I f_n.$$

□

C.2 The Weierstrass M -test

We've been developing the theory of uniform convergence for sequences of functions. Along the way, we've proved some results about the uniform convergence of series of functions, the most important of which is Corollary C.6. This corollary showed that a series is uniformly convergent if and only if it satisfies the Uniform Cauchy Criterion. As you saw in Exercise 10, while this criterion/condition is very useful, it is not terribly easy to apply. Our main result of this section, the M -test of Weierstrass, gives an relatively straightforward condition guaranteeing that a given series converges uniformly. We will then amass some facts following from this result which will be used in our study of Fourier series.

Theorem C.10 (The Weierstrass M -test). *Let $I = [a, b]$ be an interval and consider a sequence of bounded complex-valued functions $\{f_n\}$ on I . For each $n \in \mathbb{N}$, set*

$$M_n = \|f_n\|_\infty = \sup_{x \in I} |f_n(x)|.$$

If the series $\sum_{n=1}^\infty M_n$ converges, then the series $\sum_{n=1}^\infty f_n$ converges uniformly on I .

Before giving the proof, observe that the series $\sum_{n=1}^\infty M_n$ is a series of non-negative numbers and determining the convergence of this series is the subject matter of introductory calculus. This is usually an easier condition to verify than the Cauchy criterion.

Proof. We will verify that the Cauchy criterion (Corollary C.6) is satisfied for the series $\sum_n f_n$. To this end, let $\epsilon > 0$. Given that $\sum_n M_n$ converges, its partial sums are necessarily a Cauchy sequence and so there must be some natural number N for which

$$\sum_{k=n}^m M_k \leq \sum_{k=n-1}^m M_k = \sum_{k=1}^m M_k - \sum_{k=1}^{n-1} M_k = \left| \sum_{k=1}^m M_k - \sum_{k=1}^{n-1} M_k \right| < \epsilon$$

whenever $m \geq n \geq N$. Here, we have used the fact that $M_k \geq 0$ for all k . Observe now that, for any $x \in I$ and $m \geq n \geq N$, the triangle inequality guarantees that

$$\left| \sum_{k=n}^m f_k(x) \right| \leq \sum_{k=n}^m |f_k(x)| \leq \sum_{k=n}^m \|f_k\|_\infty = \sum_{k=n}^m M_k < \epsilon,$$

as desired. □

Following directly from Theorems C.10 and C.7 and Corollary C.9, we obtain the following corollary.

Corollary C.11. *Let I be an interval and let $\{f_k\}$ be a sequence of complex-valued functions on I , i.e., $\{f_k\} \subseteq C(I)$. For each $n \in \mathbb{N}$, set*

$$M_n = \|f_n\|_\infty = \sup_{x \in I} |f_n(x)|.$$

If the series $\sum_{n=1}^\infty M_n$ converges, then $\sum_{n=1}^\infty f_n$ converges uniformly on I and its sum

$$f(x) = \sum_{n=1}^\infty f_n(x)$$

is a continuous function on I , i.e., $f \in C(I)$. Further,

$$\int_I f = \int_I \sum_{n=1}^\infty f_n = \sum_{n=1}^\infty \int_I f_n.$$

Proof. The statement regarding uniform convergence follows directly from Theorem C.10. Because f_n is continuous for each n , the partial sums $\{S_n\}$ are necessarily continuous functions on I . The uniform convergence of the series is the statement that the partial sums converge uniformly to the sum of the series and so, by virtue of Theorem C.7, the sum f is necessarily continuous on I . Finally, upon noting that $\{f_n\} \subseteq C(I) \subseteq R(I)$, an appeal to Corollary C.9 gives the final statement immediately. □

Exercise 24

The Weierstrass M -test says that the “ M condition”, i.e., the condition that $\sum_{n=1}^\infty M_n$ converges, is a sufficient condition for the uniform convergence of the series $\sum f_n$. This is in contrast to Corollary C.6 which gives a condition both necessary and sufficient for uniform convergence. Show that that “ M condition” (of the Weierstrass M -test) is not necessary for convergence. That is, find a sequence of functions $\{f_n\}$ on an interval I for which $\sum_{n=1}^\infty f_n$ converges uniformly yet $\sum_{n=1}^\infty M_n = \infty$ for $M_n = \|f_n\|_\infty$. Hint: A nice example can

be produced which is an alternating series. Feel free to use results from introductory calculus (such as the alternating series test).

C.3 Defining Convergence with the Integral: A glimpse at Lebesgue norms

As the supremum norm $\|\cdot\|_\infty$ allows us to measure the “size” of a function bounded function (and with it you were able to characterize uniform convergence), the integral also allows us to measure the “size” of a function by integrating its absolute value. Measuring the size of functions with the integral turns out to be a very fruitful activity. To formalize things, I will take this opportunity to introduce a class of “norms” on functions, called the Lebesgue norms or the L^p norms, of which the supremum norm is an important example. To this end, we fix an interval I and, for each $1 \leq p < \infty$, we define the $L^p(I)$ norm of a function $f \in R(I)$ by

$$\|f\|_p = \left(\int_I |f(x)|^p dx \right)^{1/p}.$$

For $p = \infty$, we have as before

$$\|f\|_p = \|f\|_\infty = \sup_{x \in I} |f(x)|$$

for $f \in R(I)$. For each $1 \leq p \leq \infty$, each L^p norm gives us a different way to measure the “size” of a function. Let’s accumulate some facts about these norms.

Proposition C.12. *Given an interval I and $1 \leq p \leq \infty$, let $\|\cdot\|_p$ denote the $L^p(I)$ norm defined above. Then, for any $f, g \in R(I)$ and $\alpha \in \mathbb{C}$, we have*

1.

$$\|f\|_p \geq 0$$

2.

$$\|\alpha f\|_p = |\alpha| \|f\|_p$$

3.

$$\|f + g\|_p \leq \|f\|_p + \|g\|_p$$

Truthfully, the above proposition only guarantees that $\|\cdot\|_p$ is a so-called *semi-norm* on $R(I)$ because there are non-zero functions $f \in R(I)$ for which $\|f\|_p = 0$.

Proof. As you have already shown that these properties hold when $p = \infty$ (Exercise 9), we shall assume that $1 \leq p < \infty$. Now, because the integral of a non-negative function is non-negative, the validity of Item 1 is clear. Also, for $f \in R(I)$ and $\alpha \in \mathbb{C}$,

$$\|\alpha f\|_p^p = (\|\alpha f\|_p)^p = \int_I |\alpha f(x)|^p dx = \int_I |\alpha|^p |f(x)|^p dx = |\alpha|^p \int_I |f(x)|^p dx$$

from which we immediately obtain Item 2. It remains to prove Item 3, also called Minkowski’s inequality. This inequality is most easily obtained using the machinery of measure theory, though our proof here only relies on the convexity of the function $\mathbb{C} \ni z \mapsto |z|^p$, a fact which can be established using only elementary calculus.

To this end, we first assume show that, if $h_1, h_2 \in R(I)$ are such that $\|h_1\|_p, \|h_2\|_p \leq 1$, then, for any $0 \leq t \leq 1$, $\|th_1 + (1-t)h_2\|_p \leq 1$. This is equivalently the statement that the unit ball

$$B_p = \{h \in R(I) : \|h\|_p \leq 1\}$$

is a convex set. Let us fix $0 \leq t \leq 1$ and $h_1, h_2 \in B_p$ and observe that the convexity of the map $z \mapsto |z|^p$ guarantees that

$$|th_1(x) + (1-t)h_2(x)|^p \leq t|h_1(x)|^p + (1-t)|h_2(x)|^p$$

for all $x \in I$. I'll make note that the convexity used here for complex numbers is also called the supporting hyperplane property and can be understood geometrically as the graph of the function $|z|^p$ always living below its secant lines/planes. In view of this inequality, the monotonicity of the integral guarantees that

$$\int_I |th_1(x) + (1-t)h_2(x)|^p dx \leq t \int_I |h_1(x)|^p dx + (1-t) \int_I |h_2(x)|^p dx$$

or equivalently

$$\|th_1 + (1-t)h_2\|_p^p \leq t\|h_1\|_p^p + (1-t)\|h_2\|_p^p.$$

Recalling that $\|h_1\|_p \leq 1$ and $\|h_2\|_p \leq 1$, we conclude that

$$\|th_1 + (1-t)h_2\|_p^p \leq t \cdot 1 + (1-t) \cdot 1 = 1$$

and so $\|th_1 + (1-t)h_2\|_p \leq 1$, as was asserted.

We now get to the task at hand. Let $f, g \in R(I)$ and we shall assume that $\|f\|_p$ and $\|g\|_p$ are non-zero (treating these trivial cases is much more simple). We write

$$\frac{f+g}{\|f\|_p + \|g\|_p} = \frac{\|f\|_p}{\|f\|_p + \|g\|_p} \frac{f}{\|f\|_p} + \frac{\|g\|_p}{\|f\|_p + \|g\|_p} \frac{g}{\|g\|_p} = t \frac{f}{\|f\|_p} + (1-t) \frac{g}{\|g\|_p}$$

where $t = \|f\|_p / (\|f\|_p + \|g\|_p)$ is a number between 0 and 1. By virtue of Item 2, both $h_1 = f/\|f\|_p$ and $h_2 = g/\|g\|_p$ have L^p norm 1. In view of the property proved in the preceding paragraph, we conclude that

$$\left\| \frac{f+g}{\|f\|_p + \|g\|_p} \right\|_p = \|th_1 + (1-t)h_2\|_p \leq 1.$$

Therefore, a final appeal to Item 2 gives the inequality

$$\frac{1}{\|f\|_p + \|g\|_p} \|f+g\|_p \leq 1$$

from which the desired result follows without trouble. □

With these norms and this way of measuring functions, we can define new notions of convergence. To this end, given a sequence of functions $\{f_n\} \subseteq R(I)$ and $f \in R(I)$, we say that $\{f_n\}$ converges to f in $L^p(I)$ or with respect to the L^p norm if

$$\lim_{n \rightarrow \infty} \|f_n - f\|_p = 0.$$

There are three L^p norms that will be of particular interest for us, $p = 1, 2$ and ∞ . In the case that $p = 2$, there is an additional structure with which you are already familiar from linear algebra, the inner product (a generalization of the dot product). For integrable functions f and g , we define the L^2 inner product of f and g to be the number

$$\langle f, g \rangle = \int_I f(x) \overline{g(x)} dx.$$

As it is easy to verify using properties of the integral, the $L^2(I)$ inner product satisfies the following properties:

1.

$$\langle f, g \rangle = \overline{\langle g, f \rangle} \quad \text{for } f, g \in R(I)$$

2.

$$\langle \alpha f + \beta h, g \rangle = \alpha \langle f, g \rangle + \beta \langle h, g \rangle \quad \text{for } f, g, h \in R(I) \text{ and } \alpha, \beta \in \mathbb{C}.$$

3.

$$\langle g, \alpha f + \beta h \rangle = \overline{\alpha} \langle g, f \rangle + \overline{\beta} \langle g, h \rangle \quad \text{for } f, g, h \in R(I) \text{ and } \alpha, \beta \in \mathbb{C}.$$

We also notice, that the L^2 inner product recaptures the L^2 norm:

$$\|f\|_2 = \left(\int_I |f(x)|^2 dx \right)^{1/2} = \left(\int_I f(x) \overline{f(x)} dx \right)^{1/2} = \sqrt{\langle f, f \rangle}$$

for $f \in R(I)$. An extremely important property of the L^2 inner product is captured by the following theorem.

Theorem C.13 (The Cauchy-Schwarz Inequality). *For any $f, g \in R(I)$,*

$$|\langle f, g \rangle| \leq \|f\|_2 \|g\|_2$$

Proof. Let's first assume that $h_1, h_2 \in R(I)$ have $\|h_1\|_2 = \|h_2\|_2 = 1$. We observe that, for any $x \in I$,

$$0 \leq (|h_1(x)| - |h_2(x)|)^2 = (|h_1(x)|^2 + |h_2(x)|^2 - 2|h_1(x)||h_2(x)|).$$

Therefore

$$|h_1(x)||h_2(x)| \leq \frac{|h_1(x)|^2}{2} + \frac{|h_2(x)|^2}{2}$$

for all $x \in I$. By virtue of Proposition B.8, the preceding inequality shows that

$$\begin{aligned} |\langle h_1, h_2 \rangle| &= \left| \int_I h_1(x) \overline{h_2(x)} dx \right| \\ &\leq \int_I |h_1(x)||h_2(x)| dx \\ &\leq \frac{1}{2} \int_I |h_1(x)|^2 dx + \frac{1}{2} \int_I |h_2(x)|^2 dx \\ &\leq \frac{1}{2} \|h_1\|_2^2 + \frac{1}{2} \|h_2\|_2^2 = 1. \end{aligned}$$

Thus $|\langle h_1, h_2 \rangle| \leq 1$ whenever $h_1, h_2 \in R(I)$ have unit L^2 -norm. Now, given any $f, g \in R(I)$ with non-zero L^2 norms, we observe that $h_1 = f/\|f\|_2$ and $h_2 = g/\|g\|_2$ have $\|h_1\|_2 = \|h_2\|_2 = 1$ and so by the properties of the L^2 inner product outlined above

$$|\langle f, g \rangle| = \|f\|_2 \|g\|_2 \left| \left\langle \frac{f}{\|f\|_2}, \frac{g}{\|g\|_2} \right\rangle \right| = \|f\|_2 \|g\|_2 |\langle h_1, h_2 \rangle| \leq \|f\|_2 \|g\|_2$$

as desired.

Finally, let us assume that $\|f\|_2 = 0$ or $\|g\|_2 = 0$. In this final case, our job is to show that $\langle f, g \rangle = 0$ because the right-hand side of the Cauchy-Schwarz inequality is zero. Without loss of generality we assume that $\|g\|_2 = 0$ and observe that, for all $t \in \mathbb{R}$,

$$\begin{aligned} \|f + tg\|_2^2 &= \langle f + tg, f + tg \rangle = \langle f, f \rangle + \langle f, tg \rangle + \langle tg, f \rangle + \langle tg, tg \rangle \\ &= \|f\|_2^2 + \langle f, tg \rangle + \overline{\langle f, tg \rangle} + t^2 \|g\|_2^2 \\ &= \|f\|_2^2 + 2 \operatorname{Re}(\langle f, tg \rangle) + 0 \\ &= \|f\|_2^2 + 2t \operatorname{Re}(\langle f, g \rangle) \end{aligned}$$

where we have used the fact that t is real and $z + \bar{z} = 2 \operatorname{Re} z$ for any complex number z (this is something you should check). In view of the equation above, we have

$$0 \leq \|f\|_2^2 + 2t \operatorname{Re}(\langle f, g \rangle)$$

for all $t \in \mathbb{R}$. I claim that this inequality implies that $\operatorname{Re}(\langle f, g \rangle) = 0$. If $\operatorname{Re}(\langle f, g \rangle) \neq 0$, then setting $t = -(\|f\|_2^2 + 1)/\operatorname{Re}(\langle f, g \rangle)$ in the above inequality yields

$$0 \leq \|f\|_2^2 + 2 \left(-\frac{\|f\|_2^2 + 1}{\operatorname{Re}(\langle f, g \rangle)} \right) \operatorname{Re}(\langle f, g \rangle) = \|f\|_2^2 - 2\|f\|_2^2 - 2 = -(\|f\|_2^2 + 2)$$

which is impossible because $\|f\|_2^2 + 2 \geq 2 > 0$. From this we conclude that $\operatorname{Re}(\langle f, g \rangle) = 0$. An analogous argument (done by expanding $\|f + itg\|_2^2$) shows that $\operatorname{Im}(\langle f, g \rangle) = 0$. All together, we conclude that $\langle f, g \rangle = 0$. \square

There are many generalizations of the Cauchy-Schwarz inequality that turn out to be useful for Fourier analysis. The following one, which we give without proof, is called Hölder’s inequality [?]. The theorem essentially says that the integral of a product of functions f and g is bounded above in absolute value by the L^p norm of f and the L^q norm of g where $1 \leq p, q \leq \infty$ are such that

$$\frac{1}{p} + \frac{1}{q} = 1.$$

Such a pair p and q are said to be conjugate exponents and here we assume the convention that $1/\infty = 0$. So, for example $p = 2$ and $q = 2$ are conjugate exponents. Also $p = 1$ and $q = \infty$ are conjugate exponents.

Theorem C.14 (Hölder’s inequality). *Let $1 \leq p, q \leq \infty$ be conjugate exponents. Then, for any $f, g \in R(I)$, the product fg is integrable and*

$$\left| \int_I f(x)g(x) dx \right| \leq \|f\|_p \|g\|_q.$$

Exercise 25

Though we’ve already proven the triangle inequality for the L^p norm (also called the Minkowski inequality), please show that the triangle inequality

$$\|f + g\|_p \leq \|f\|_p + \|g\|_p$$

is a consequence of Hölder’s inequality (and thus the latter is more “fundamental”). Hint: First observe that $|f(x) + g(x)|^p \leq |f(x) + g(x)|^{p-1}(|f(x)| + |g(x)|)$ for all x . Then apply Hölder’s inequality to the terms on the right-hand side.

As an application of Hölder’s inequality, we have the following theorem which gives a relationship to convergence between L^p norms.

Theorem C.15. *Let $I = [a, b]$ be a bounded interval and let $\{f_n\}$ be a sequence in $R(I)$. Also, let $f \in R(I)$. Given any $1 \leq r \leq s \leq \infty$, if*

$$\lim_{n \rightarrow \infty} \|f_n - f\|_s = 0 \quad \text{then} \quad \lim_{n \rightarrow \infty} \|f_n - f\|_r = 0.$$

If you take a course in measure theory, you will learn that this result depends critically on the fact that $I = [a, b]$ is a bounded interval. Before giving the proof (taking Hölder’s inequality for granted), we note that it implies the following statement (as a special case).

$$\text{If } \lim_{n \rightarrow \infty} \|f_n - f\|_\infty = 0 \quad \text{then} \quad \lim_{n \rightarrow \infty} \|f_n - f\|_1 = \lim_{n \rightarrow \infty} \int_I |f_n(x) - f(x)| dx = 0.$$

This statement should be familiar as it recaptures Theorem C.8 in view of the correspondence between uniform convergence and convergence in the L^∞ norm. Now let’s prove the theorem.

Proof. Fixing $1 \leq r \leq s$, set $p = s/r$ and observe that $p \geq 1$. In the case that $r = s = \infty$, the assertion is obvious. We therefore assume that $r < \infty$ and, in view of Hölder’s inequality, we obtain

$$\|f_n - f\|_r^r = \int_I |f_n(x) - f(x)|^r dx = \int_I |f_n(x) - f(x)|^r \cdot 1 dx \leq \|(f_n - f)^r\|_p \|1\|_q \tag{7}$$

where q is the conjugate exponent to p and 1 is the constant function. If $p = \infty$, necessarily $s = \infty$, $q = 1$ and we have

$$\|(f_n - f)^r\|_p = \sup_{x \in I} |f_n(x) - f(x)|^r = \|f_n - f\|_\infty^r. \tag{8}$$

In this case, combining the two preceding inequalities guarantee that

$$\|f_n - f\|_r^r \leq \|f_n - f\|_\infty^r \|1\|_1 = \|f_n - f\|_\infty^r |b - a|$$

or, equivalently,

$$\|f_n - f\|_r \leq (b - a)^{1/r} \|f_n - f\|_\infty.$$

If $p < \infty$, we note that

$$\begin{aligned} \|(f_n - f)^r\|_p &= \left(\int_I (|f_n(x) - f(x)|^r)^p dx \right) \\ &= \left(\int_I |f_n(x) - f(x)|^{pr} dx \right)^{1/p} \\ &= (\|f_n - f\|_s^s)^{1/p} = \|f_n - f\|_s^{s/p} = \|f_n - f\|_s^r \end{aligned}$$

where we have used the fact that $pr = s$ and $s/p = r$. Combining this with (7) yields

$$\|f_n - f\|_r^r \leq \|f_n - f\|_s^r \|1\|_q = \|f_n - f\|_s^r \|1\|_q$$

and therefore

$$\|f_n - f\|_r \leq \|f_n - f\|_s \|1\|_q^{1/r}.$$

Finally, noting that

$$\|1\|_q = \begin{cases} (\int_I 1^q)^{1/q} = (b - a)^{1/q} & q < \infty \\ 1 & q = \infty \end{cases} = (b - a)^{1/q}$$

(as long as we interpret $1/\infty = 0$, we have

$$\|f_n - f\|_r \leq \|f_n - f\|_s (b - a)^{1/rq} = (b - a)^{(\frac{1}{r} - \frac{1}{s})} \|f_n - f\|_s \tag{9}$$

where we have used the fact that $\frac{1}{r} = \frac{1}{rp} + \frac{1}{rq} = \frac{1}{s} + \frac{1}{rq}$. Combining both cases (7) and (9) (and using the conventions that $1/0 = \infty$ and $1/\infty = 0$, we obtain

$$\|f_n - f\|_r \leq (b - a)^{(\frac{1}{r} - \frac{1}{s})} \|f_n - f\|_s$$

whenever $1 \leq r \leq s$. Finally, if the sequence $\{f_n\}$ has $\lim_{n \rightarrow \infty} \|f_n - f\|_s = 0$, the preceding inequality guarantees that $\lim_{n \rightarrow \infty} \|f_n - f\|_r = 0$. □

Example 4

To illustrate the preceding theorem, let's construct a sequence of functions which converge to the zero function with respect to the L^s norm for "small" s but diverge in the L^s norm for "large" s . To this end, set $I = [-1, 1]$ and fix $0 < a \leq \infty$. For each $n \in \mathbb{N}$, define

$$f_n(x) = n^{1/a} e^{-n|x|} \quad \text{for } -1 \leq x \leq 1.$$

We are assuming the convention that $n^{1/a} = n^0 = 1$ when $a = \infty$. Figure 12 illustrates f_2 and f_{10} in the case that $a = 1$.




Figure 12: The graphs of f_2 and f_{10} when $a = 1$.

A study of this particular sequence of functions provides a nice way to understand which factors contribute to the L^s norm of a function. For this sequence f_n , for a value of $a < \infty$, we see that the peaks at $f_n(x)$ (which happen at $x = 0$) grow unboundedly while the graphs become more and more narrow as $n \rightarrow \infty$. In terms of area under the graph, which is the essential contributor to the L^s norms, this can be seen as a competition between growing height and shrinking width. Let's nail things down precisely.

As suggested by the figure, it is easily verified that, for each n , f_n is continuous on the interval I , i.e., $\{f_n\} \subseteq C(I)$, and therefore $\{f_n\}$ is a sequence of Riemann integrable functions. Let's compute the $L^s(I)$ norms of this sequence: For $s = \infty$, we have

$$\|f_n\|_s = \|f_n\|_\infty = \sup_{x \in I} |f_n(x)| = n^{1/a}.$$

for each $n \in \mathbb{N}$. For $1 \leq s < \infty$, we have

$$\begin{aligned} \|f_n\|_s &= \left(\int_I |f_n(x)|^s dx \right)^{1/s} \\ &= \left(\int_{-1}^1 n^{s/a} e^{-sn|x|} dx \right)^{1/s} \\ &= n^{1/a} \left(2 \int_0^1 e^{-snx} dx \right)^{1/s} \\ &= n^{1/a} 2^{1/s} \left(\frac{e^{-snx}}{-sn} \Big|_{x=0}^{x=1} \right)^{1/s} \\ &= n^{1/a} \left(\frac{2}{sn} \right)^{1/s} (1 - e^{-sn})^{1/s} \\ &= n^{(1/a - 1/s)} \left(\frac{2}{s} \right)^{1/s} \left(1 - \frac{1}{e^{sn}} \right)^{1/s} \end{aligned}$$

for each $n \in \mathbb{N}$. We therefore have the following behavior: if $s < a$, then $1/a - 1/s < 0$ (where we can't have $s = \infty$) and so

$$\lim_{n \rightarrow \infty} \|f_n - 0\|_s = \lim_{n \rightarrow \infty} \|f_n\|_s = \lim_{n \rightarrow \infty} n^{1/a - 1/s} (2/s)^{1/s} (1 - 1/e^{sn})^{1/s} = 0 \cdot (2/s)^{1/s} \cdot 1 = 0.$$

Consequently, if $s < a$, $\{f_n\}$ converges to the zero function with respect to the $L^s(I)$ norm. If $s \geq a$, then, for $s = \infty$,

$$\lim_{n \rightarrow \infty} \|f_n - 0\|_s = \lim_{n \rightarrow \infty} n^{1/a} = \infty$$

and, for $s < \infty$ $1/a - 1/s \geq 0$,

$$\lim_{n \rightarrow \infty} \|f_n - 0\|_s = \lim_{n \rightarrow \infty} n^{(1/a-1/s)}(2/s)^{1/s}(1 - 1/e^{sn})^{1/s} = \begin{cases} \infty & a < s \\ (2/s)^{1/s} & a = s. \end{cases}$$

In other words, the sequence $\{f_n\}$ converges to 0 for all $s < a$ (all small s) and does not converge to 0 for all $s \geq a$ (all large s). In particular, upon fixing $s < a$, if $r \leq s$, then $\{f_n\}$ converges to zero in both L^s and L^r norms. If $r > s$, then it is possible to $\{f_n\}$ to not converge to zero in the L^r norm (namely, when $r \geq a$) while still converging to zero in the L^s norm. As it must be, this is consistent with the preceding theorem.