



## APRIL 26 LECTURE

### SUPPLEMENTARY REFERENCES:

- *Single Variable Calculus*, Stewart, 7th Ed.: Section 8.1.
- - *Integral Calculus*, Khan Academy: Area & arc length using calculus.

KEYWORDS: arc length

---

## APPLICATIONS OF INTEGRATION: ARC LENGTH

In this lecture we will investigate the notion of *arc length*. We define what we mean by the *length* of the graph of a continuous function  $f(x)$ , and deduce an integration formula for the arc length of a large class of curves in the plane.

### Lines and length

Given a curve  $C$  in the plane, the notion of what we mean by its length seems intuitive: take a piece of string, or some other pliable material, lay the string along the curve and mark the endpoints. After pulling the string taut we can measure the length of the curve - ta-da!




But how do we perform this procedure **in practice**? Let's start out simple and look at straight line segments.


MATHEMATICAL WORKOUT - FLEX THOSE MUSCLES!

1. Find the length of the straight line segment from  $(1, 2)$  to  $(5, 4)$ . (*Hint: realise the straight line segment as the hypotenuse of a right triangle*)

2. Generalise the above example to find a formula for the length of the segment of the straight line  $y = mx + c$  from  $x = a$  to  $x = b$ .




3. What's the length of the following curve? Each straight line segment has slope  $m_1, m_2, m_3, m_4$ .




### The arc length formula

Consider the function  $f(x) = 4 - x^2$ . How might we determine the length of the parabolic curve segment between  $x = -2$  and  $x = 2$ ?



CHECK YOUR UNDERSTANDING

Subdivide the interval  $[-2, 2]$  into four segments of equal length as demonstrated below



1. Write down the slopes  $m_1$ ,  $m_2$ ,  $m_3$  and  $m_4$  of the line segments drawn above

$m_1 =$ _____	$m_2 =$ _____	$m_3 =$ _____	$m_4 =$ _____
---------------	---------------	---------------	---------------

2. Determine the total length of curve consisting of the three dashed straight line segments above. How is this distance related to the actual length of the parabolic curve segment?

3. Determine  $x_1, x_2, x_3, x_4$  in the interval  $[-2, 2]$  such that

$$f'(x_1) = m_1, \quad f'(x_2) = m_2, \quad f'(x_3) = m_3, \quad f'(x_4) = m_4.$$

4. How can you interpret the previous problem using the graph of  $f(x)$ ?

Let's consider what happens if we subdivide the interval  $[-2, 2]$  into  $n$  segments having equal length and look to *approximate* the length  $L$  of the parabolic curve segment using a piecewise linear curve  $C_n$  (i.e. a collection of  $n$  straight line segments, analogous to what we did above).

1. First we observe that each segment will have length \_\_\_\_\_.
2. If each straight line segment in  $C_n$  has slope  $m_1, m_2, \dots, m_n$ , respectively, then

Length of  $C_n =$  \_\_\_\_\_

3. In the  $j^{\text{th}}$  segment of  $[-2, 2]$ , we can find  $x_j$  such that  $f'(x_j) =$  \_\_\_\_\_. Hence, we can rewrite

Length of  $C_n =$  \_\_\_\_\_

Now, as  $n$  gets very large, we expect that the length of  $C_n$  will give a good approximation of the length of the parabolic curve segment from  $x = -2$  to  $x = 2$ . It seems reasonable, therefore, that we should be able to write

$$L = \lim_{n \rightarrow \infty} \text{Length of } C_n = \lim_{n \rightarrow \infty} \text{_____}$$

We recognise this last expression as the Riemann sum associated to the function

$$g(x) = \text{_____}$$

Let  $f(x)$  be a function defined on the interval  $[a, b]$  for which the derivative  $f'(x)$  is continuous (this will be the case for all functions we encounter). Then, we define the **arc length of the curve  $y = f(x)$  between  $x = a$  to  $x = b$**  to be

\_\_\_\_\_

**Remark:** The condition that  $f'(x)$  is continuous is required to ensure that the integral (i.e. a limit of Riemann sums) is well-defined.

In order to determine definite integrals, we need to recall the following essential result.

## Fundamental Theorem of Calculus, Revisited


Let  $F(x)$  be an antiderivative of  $f(x)$ . Then,

$$\int_a^b f(x)dx = F(b) - F(a)$$

### Example:

1. The arc length of the curve  $y = \frac{2}{3}x^{\frac{3}{2}}$  between  $x = 0$  and  $x = 1$  is given by

$$\int_0^1 \sqrt{1+x} dx \stackrel{u=1+x}{=} \left[ \frac{2}{3}(1+x)^{\frac{3}{2}} \right]_0^1 = \frac{2\sqrt{8}}{3} - \frac{2}{3}$$



2. The arc length of the curve  $y = 4 - x^2$  between  $x = -2$  and  $x = 2$  is given by

$$\int_{-2}^2 \sqrt{1+4x^2} dx.$$

To determine this integral we will need to make an inverse trigonometric substitution  $x = \frac{1}{2} \tan(t)$ . This leads to the definite integral

$$2 \int_0^{\arctan(2)} \sec^3(t) dt$$

Using integration by parts

$$\begin{array}{ll} f = \sec(t) & g' = \sec^2(t) \\ f' = \sec(t) \tan(t) & g = \underline{\hspace{2cm}} \end{array}$$

and

$$\int \sec^3(t) dt = \sec(t) \tan(t) - \int \sec(t) \tan^2(t) dt = \sec(t) \tan(t) - \underline{\hspace{4cm}}$$

$$\implies \int \sec^3(t) dt = \underline{\hspace{4cm}}$$

3. Let  $f(x) = \frac{x^3}{6} + \frac{1}{2x}$ , so that  $f'(x) = \frac{x^2}{2} - \frac{1}{2x^2}$ . The arc length of the curve  $y = f(x)$  between  $x = 1$  and  $x = 3$  is given by

$$\begin{aligned}\int_1^3 \sqrt{1 + \frac{1}{4}(x^2 - x^{-2})^2} dx &= \int_1^3 \sqrt{x^2 + \frac{1}{2} + x^{-2}} dx \\ &= \int_1^3 \sqrt{\frac{1}{4}(x^2 + x^{-2})^2} dx \\ &= \frac{1}{2} \int_1^3 (x^2 + x^{-2}) dx, \quad \text{since } x^2 + x^{-2} \geq 0 \text{ whenever } 1 \leq x \leq 3, \\ &= \frac{1}{2} \left[ \frac{x^3}{3} - \frac{1}{x} \right]_1^3 = \frac{14}{3}\end{aligned}$$

CHECK YOUR UNDERSTANDING

Determine the arc length of  $y = 2(x - 1)^{3/2}$  between  $x = 1$  and  $x = 5$ .