



## APRIL 30 LECTURE

### SUPPLEMENTARY REFERENCES:

- *Single Variable Calculus*, Stewart, 7th Ed.: Section 8.1.
- *Integral Calculus*, Khan Academy: Area & arc length using calculus.

KEYWORDS: surface area, surface of revolution


## APPLICATIONS OF INTEGRATION: SURFACE AREA & SURFACES OF REVOLUTION

In the last lecture I was **frustrated by the frustrum!** Let's sort things out.


We are trying to determine the surface area of the general frustrum  $F$  obtained by rotating a line segment  $y = mx + c$ ,  $a \leq x \leq b$ , around the  $x$ -axis.

$$y = mx + c$$

$$a \leq x \leq b$$



To determine the surface area of  $F$  we cut the frustrum along a line and unroll it in the plane we obtain a circular sector having radius  $r$  and angle  $\theta$  with a concentric sector of radius  $r - s$  removed.



We have

$$r_1 = ma + c, \quad r_2 = mb + c$$

and using the formula for the length of the line segment  $y = mx + c$ ,  $x_1 \leq x \leq x_2$ ,

$$(x_2 - x_1)\sqrt{1 + m^2}$$

we determine

$$s = \left(a + \frac{c}{m}\right) \sqrt{1 + m^2}, \quad r = \left(b + \frac{c}{m}\right) \sqrt{1 + m^2}$$

Moreover, since  $p = r - s$ , we obtain

$$\theta p = \theta(r - s) = 2\pi(r_2 - r_1) = 2\pi m(b - a)$$

Here we use the fact that  $\theta s = 2\pi r_1$  (resp.  $\theta r = 2\pi r_2$ ) is the length of a circular arc appearing above. Now, the area of  $F$  is given by

$$\begin{aligned} A &= \frac{\theta}{2\pi} \pi(r^2 - s^2) = \frac{\theta}{2} p(r + s) = \pi m(b - a) \sqrt{1 + m^2} \left(a + b + \frac{2c}{m}\right) \\ \implies A &= 2\pi \sqrt{1 + m^2} \left(\frac{m}{2}(b^2 - a^2) + c(b - a)\right) \end{aligned} \quad (*)$$

**PHEW!**

CHECK YOUR UNDERSTANDING

Let  $f(x) = mx + c$ . Show that  $A$  given in (\*) can be computed using a definite integral:

$$A = 2\pi \int_a^b f(x) \sqrt{1 + f'(x)^2} dx$$

We can now approximate the surface area of a surface of revolution using a collection of circular frustrums.

PICTURE

Let  $S$  be a surface of revolution obtained from  $f(x)$ ,  $a \leq x \leq b$ . Choose a natural number  $n$ .

1. Subdivide  $[a, b]$  into  $n$  subintervals having equal length so that the endpoints of each subinterval are

$$a = x_0 < x_1 < x_2 < \dots < x_n = b$$

2. Define the piecewise linear function  $g_n(x)$  as follows (it's the function whose graph is the collection of line segments above):

$$g_n(x) = m_i(x - x_i) + f(x_i), \quad \text{when } x_{i-1} \leq x \leq x_i.$$

Here

$$m_i = \underline{\hspace{2cm}}$$

The piecewise linear function  $g_n(x)$  provides an approximation to the graph of  $f(x)$

3. Then, the surface of revolution  $S$  is approximated by a collection of  $n$  circular frustrums  $F_1, \dots, F_n$  as in the above diagram. Moreover,

Surface area of  $F_i = \underline{\hspace{3cm}}$

4. Hence, the surface area of  $S$  is obtained as the limit

$$\begin{aligned} \text{Surface area of } S &= \lim_{n \rightarrow \infty} \sum_{i=1}^n 2\pi \int_{x_{i-1}}^{x_i} \sqrt{1 + (g'_n(x))^2} g_n(x) dx \\ &= \underline{\hspace{3cm}} \end{aligned}$$

**Example:** The surface area  $A$  of the surface of revolution about the  $x$ -axis obtained from  $f(x) = 2\sqrt{x}$  when  $1 \leq x \leq 2$  is

$$A = 2\pi \int_1^2 \sqrt{1 + \frac{1}{x}} 2\sqrt{x} dx = 4\pi \int_1^2 \sqrt{x+1} dx = 4\pi \left[ \frac{2}{3}(x+1)^{3/2} \right]_1^2 = \frac{8\pi}{3} (\sqrt{27} - \sqrt{8})$$

CHECK YOUR UNDERSTANDING

The surface of the ball of radius  $a > 0$  can be realised as a surface of revolution of the function  $f(x) = \sqrt{a^2 - x^2}$ ,  $-a \leq x \leq a$ .

1. Show that

$$1 + f'(x)^2 = \frac{a^2}{a^2 - x^2}$$

2. Show that

$$f(x)\sqrt{1 + f'(x)^2} = a$$

3. Use the formula for surface area of a surface of revolution and deduce the well-known formula for the surface area  $A$  of a ball of radius  $a$ :

$$A = 4\pi a^2$$