



MARCH 22 LECTURE

SUPPLEMENTARY REFERENCES:

- *Single Variable Calculus*, Stewart, 7th Ed.: Section 6.1, 6.2*, 6.3*, 6.6.
- *Calculus*, Spivak, 3rd Ed.: Section 18.

KEYWORDS: chain rule, natural logarithm

THE NATURAL LOGARITHM II & INVERSE TRIGONOMETRIC FUNCTIONS

We saw yesterday that the exponential function $\exp(x)$ admits an inverse function $\exp^{-1}(x)$. We also determined that

$$\frac{d}{dx} \exp^{-1}(x) = \frac{1}{x} \quad (*)$$

A Fundamental Interlude

Let $f(x)$ be a function. An **antiderivative** of $f(x)$ is a differentiable function $F(x)$ satisfying

$$\frac{d}{dx} F(x) = f(x).$$

Proposition: If $F(x)$ and $G(x)$ are antiderivatives of $f(x)$ then

$$F(x) = G(x) + c,$$

for some constant c .

The most important Theorem you saw in Calculus I was an approach to determining the antiderivative of a continuous function.

Fundamental Theorem of Calculus

Let $f(x)$ be a continuous function defined on the closed interval $a \leq x \leq b$. Then, the function

$$F(x) = \int_a^x f(u) du$$

is an antiderivative of $f(x)$.

The natural logarithm

We can restate (*) as follows:

$$\exp^{-1}(x) \text{ is an antiderivative of } g(x) = \frac{1}{x}.$$

We define the **natural logarithm function** to be the function

$$\ln(x) = \int_1^x \frac{dt}{t} \quad x > 0$$

By the Fundamental Theorem of Calculus, $\ln(x)$ is an antiderivative of $g(x) = \frac{1}{x}$. Hence, the Proposition implies that there is a constant c so that

$$\exp^{-1}(x) = \underline{\hspace{2cm}}.$$

Since $\exp(0) = \underline{\hspace{1cm}}$, we have $\underline{\hspace{2cm}}$. Hence,

$$\underline{\hspace{10cm}}.$$

Hence,

The **natural logarithm function** $\ln(x)$ defined above is the inverse function $\exp^{-1}(x)$,

$$\ln(x) = \exp^{-1}(x)$$

Remark:

1. You will show in Homework that the function $\ln(x)$ just defined as the inverse of $\exp(x)$ satisfies the expected *logarithm rules*:

- $\ln(xy) = \ln(x) + \ln(y)$, for every $x, y > 0$.
- $\ln(x^n) = n \ln(x)$, for every $x > 0$ and natural number n .

2. As the inverse function of $\exp(x)$, the following functional relationships hold:

- $\exp(\ln(y)) = y$, for every $y > 0$,
- $\ln(\exp(x)) = x$, for every x .

3. We have claimed that

$$\exp(x) = e^x, \quad \text{where } e = \exp(1) \text{ is Euler's number.}$$


Hence, we see that $\ln(x) = \log_e(x)$ is the *logarithm base e* function.

Inverse Trigonometric Functions

We are going to provide a similar analysis to determine **inverse trigonometric functions**.

CHECK YOUR UNDERSTANDING

1. Let $f(x) = \sin(x)$. Draw the graph of $f(x)$.



2. Explain why $f(x)$ is not one-to-one.


(Recall: $f(x)$ is one-to-one if *distinct inputs give distinct outputs*)

3. Determine a domain $A : a \leq x \leq b$ on which $f(x)$ is one-to-one.

4. What is the range B of $f(x)$ when the inputs are restricted to A ?

5. Explain why an inverse function $f^{-1}(y)$ to $f(x)$ exists, when we restrict to domain A .

6. Draw the graph of $f^{-1}(y)$



Definition: Consider the function $f(x) = \sin(x)$, with domain $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$. Then, $f(x)$ is one-to-one and we call its inverse function $f^{-1}(y)$ the **inverse sine function**, which we denote $\arcsin(y)$.

CHECK YOUR UNDERSTANDING

Complete the following statement:

- the domain of $\arcsin(y)$ is _____
- the range of $\arcsin(y)$ is _____

Remark:


1. We write $\arcsin(y)$ instead of $\sin^{-1}(y)$ to avoid confusion with the common notation $\sin^k(x) = (\sin(x))^k$.
2. As the inverse function of $\sin(x)$, the following functional relationship holds:

- $\sin(\arcsin(y)) = y$, for every _____,
- $\arcsin(\sin(x)) = x$, for every _____.

3. In words:

“ $\arcsin(y)$ is the arc whose sine is y ”

This is demonstrated by the following diagram: (recall that, the length of the arc drawn below is x , whenever the angle x is measured in radians)



Since $f(x) = \sin(x)$ is a differentiable function the same is true of $\arcsin(y)$. Using the formula for the derivative of an inverse function

$$\frac{d}{dy} f^{-1}(y) = \frac{1}{f'(f^{-1}(y))}$$

we have

$$\frac{d}{dy} \arcsin(y) = \underline{\hspace{10cm}}$$

Here we have used that the derivative of \sin is \cos , and used the above triangle to show that $\cos(\arcsin(y)) = \underline{\hspace{10cm}}$. Hence,

$\arcsin(x)$ is an antiderivative of $\underline{\hspace{10cm}}$

Exercise: determine the derivative of $\arctan(x)$ following a similar approach as above.